

This text is meant purely as a documentation tool and has no legal effect. The Union's institutions do not assume any liability for its contents. The authentic versions of the relevant acts, including their preambles, are those published in the Official Journal of the European Union and available in EUR-Lex. Those official texts are directly accessible through the links embedded in this document

► **B**

COMMISSION REGULATION (EC) No 943/2005
of 21 June 2005
concerning the permanent authorisation of additives in feedingstuffs
(Text with EEA relevance)
(OJ L 159, 22.6.2005, p. 6)

Amended by:

		Official Journal		
		No	page	date
► <u>M1</u>	Commission Implementing Regulation (EU) No 361/2011 of 13 April 2011	L 100	22	14.4.2011
► <u>M2</u>	Commission Implementing Regulation (EU) No 290/2014 of 21 March 2014	L 87	84	22.3.2014
► <u>M3</u>	Commission Implementing Regulation (EU) 2017/1145 of 8 June 2017	L 166	1	29.6.2017

▼B

COMMISSION REGULATION (EC) No 943/2005

of 21 June 2005

concerning the permanent authorisation of additives in feedingstuffs

(Text with EEA relevance)

Article 1

The preparation belonging to the group ‘Micro-organisms’, as specified in Annex I, is authorised for use without a time-limit as an additive in animal nutrition under the conditions laid down in that Annex.

Article 2

The preparations belonging to the group ‘Enzymes’, as specified in Annex II, are authorised for use without a time-limit as additives in animal nutrition under the conditions laid down in that Annex.

Article 3

This Regulation shall enter into force on the third day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

ANNEX I

EC No	Additive	Chemical formula, description	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
					CFU/kg of complete feeding-stuff			
Micro-organisms								
E 1705	<i>Enterococcus faecium</i> NCIMB 10415	Preparation of <i>Enterococcus faecium</i> containing a minimum of: Microencapsulated form: $1,0 \times 10^{10}$ CFU/g of additive; Granulated form: $3,5 \times 10^{10}$ CFU/g additive	Pigs for fattening	—	$0,35 \times 10^9$	$1,0 \times 10^9$	In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability to pelleting.	Without a time limit

ANNEX II

EC No	Additive	Chemical formula, description	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
					Units of activity/kg of complete feedingstuff			
Enzymes								
E 1604	► M2 ◀	Preparation of endo-1,3(4)-beta-glucanase and endo-1,4-beta-xylanase produced by <i>Penicillium funiculosum</i> (IMI SD 101) having a minimum activity of: Powder form: endo-1,3(4)-beta-glucanase: 2 000 U ⁽¹⁾ /g endo-1,4-beta-xylanase: 1 400 U ⁽²⁾ /g Liquid form: endo-1,3(4)-beta-glucanase: 500 U/ml endo-1,4-beta-xylanase: 350 U/ml	Laying hens	—	Endo-1,3(4)-beta-glucanase: 100 U Endo-1,4-beta-xylanase: 70 U	—	1. In the directions for use of the additive and premixture, indicate the storage temperature, storage life, and stability to pelleting. 2. Recommended dose per kg of complete feedingstuff: endo-1,3(4)-beta-glucanase: 100 U endo-1,4-beta-xylanase: 70 U 3. For use in compound feed rich in non-starch polysaccharides, (mainly beta-glucans and arabinoxylans), e.g. containing more than 60 % barley or 30 % wheat.	Without time-limit a
			Turkeys for fattening	—	Endo-1,3(4)-beta-glucanase: 100 U Endo-1,4-beta-xylanase: 70 U	—	1. In the directions for use of the additive and premixture, indicate the storage temperature, storage life, and stability to pelleting. 2. Recommended dose per kg of complete feedingstuff: endo-1,3(4)-beta-glucanase: 100 U endo-1,4-beta-xylanase: 70 U 3. For use in compound feed rich in non-starch polysaccharides, (mainly beta-glucans and arabinoxylans), e.g. containing more than 30 % barley or 20 % wheat.	Without time-limit a

▼ **B**

EC No	Additive	Chemical formula, description	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
					Units of activity/kg of complete feedingstuff			
E 1613	Endo-1,4-beta-xylanase EC 3.2.1.8	Preparation of endo-1,4-beta-xylanase produced by <i>Trichoderma longibrachiatum</i> (CNCM MA 6-10 W) having a minimum activity of: Powder form: 70 000 IFP ⁽³⁾ /g Powder form: 7 000 IFP/ml	Turkeys for fattening	—	1 400 IFP	—	1. In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability to pelleting. 2. Recommended dose per kg of complete feedingstuff: 1 400 IFP 3. For use in compound feed rich in non-starch polysaccharides (mainly arabinoxylans), e.g. containing more than 38 % wheat.	Without a time-limit

▼ **M3**

▼ **B**

- (1) 1 U is the amount of enzyme which liberates 5,55 micromoles of reducing sugars (maltose equivalents) from barley beta-glucan per minute at pH 5,0 and 50 °C.
 (2) 1 U is the amount of enzyme which liberates 4,00 micromoles of reducing sugars (maltose equivalents) from birchwood xylan per minute at pH 5,5 and 50 °C.
 (3) 1 IFP is the amount of enzyme which liberates 1 micromole of reducing sugars (xylose equivalents) from oat xylan per minute at pH 4,8 and 50 °C.
 (4) ► **M3** _____ ◀
 (5) ► **M3** _____ ◀
 (6) ► **M3** _____ ◀
 (7) ► **M3** _____ ◀
 (8) ► **M3** _____ ◀