

1.3.2024

**COMMISSION IMPLEMENTING REGULATION (EU) 2020/1762****of 25 November 2020**

concerning the authorisation of a preparation of *Bacillus subtilis* DSM 32324, *Bacillus subtilis* DSM 32325 and *Bacillus amyloliquefaciens* DSM 25840 as a feed additive for all poultry species for fattening or reared for laying or reared for breeding (holder of authorisation Chr. Hansen A/S)

(Text with EEA relevance)

Article 1

The preparation specified in the Annex, belonging to the additive category 'zootechnical additives' and to the functional group 'gut flora stabilisers', is authorised as an additive in animal nutrition, subject to the conditions laid down in that Annex.

Article 2

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

ANNEX

Identification number of the additive	Name of the holder of authorisation	Additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Minimum content	Maximum content	Other provisions	End of period of authorisation
						CFU/kg of complete feedingstuff with a moisture content of 12 %		CFU/l of water for drinking			
Category of zootechnical additives. Functional group: gut flora stabilisers											
4b1894	Chr. Hansen A/S	<i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus amyloliquefaciens</i> DSM 25840	<i>Additive composition</i> Preparation of <i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus amyloliquefaciens</i> DSM 25840 containing a minimum of: $3,2 \times 10^9$ CFU/g additive ($1,6 \times 10^9$ CFU <i>B. subtilis</i> DSM 32324/g; $1,0 \times 10^9$ CFU <i>B. subtilis</i> DSM 32325/g and $0,6 \times 10^9$ CFU <i>B. amyloliquefaciens</i> DSM 25840/g)	All poultry species for fattening or reared for laying or reared for breeding	—	$1,6 \times 10^9$	—	$5,4 \times 10^8$	—	1. In the directions for use of the additive and premixtures, the storage conditions and stability to heat treatment shall be indicated. 2. For use of the additive in water for drinking the homogenous dispersion of the additive shall be ensured. ► M1 3. The additive may be used simultaneously with the following coccidiostats, in accordance with their respective conditions of authorisation as feed additives: diclazuril, decoquinone, halofuginone, monensin, salinomycin, narasin, a combination of nicarbazin and narasin, and lasalocid. ◀ 4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from its use. Where those risks cannot be	16.12.2030
			<i>Characterisation of the active substance</i> Viable spores of cells of <i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus amyloliquefaciens</i> DSM 25840								
			<i>Analytical method</i> ⁽¹⁾ Enumeration in the feed additive, premixtures, feedingstuffs and water: Spread plate method on tryptone soya agar (EN 15784). Identification: Pulsed Field Gel Electrophoresis (PFGE) method.								

▼B

Identification number of the additive	Name of the holder of authorisation	Additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Minimum content	Maximum content	Other provisions	End of period of authorisation
						CFU/kg of complete feedingstuff with a moisture content of 12 %		CFU/l of water for drinking			
										eliminated or reduced to a minimum by such procedures and measures, the additive and premixtures shall be used with personal protective equipment, including breathing protection, glasses and gloves.	

(1) Details of the analytical methods are available at the following address of the Reference Laboratory: <https://ec.europa.eu/jrc/en/eurl/feed-additives/evaluation-reports>