

Official Journal

of the European Communities

ISSN 0378-6978

L 339

Volume 39

30 December 1996

English edition

Legislation

Contents

I *Acts whose publication is obligatory*

- ★ Commission Directive 96/77/EC of 2 December 1996 laying down specific purity criteria on food additives other than colours and sweeteners⁽¹⁾ 1

Price: ECU 19,50

⁽¹⁾ Text with EEA relevance.

EN

Acts whose titles are printed in light type are those relating to day-to-day management of agricultural matters, and are generally valid for a limited period.

The titles of all other Acts are printed in bold type and preceded by an asterisk.

I

(Acts whose publication is obligatory)

COMMISSION DIRECTIVE 96/77/EC

of 2 December 1996

laying down specific purity criteria on food additives other than colours and sweeteners

(Text with EEA relevance)

THE COMMISSION OF THE EUROPEAN COMMUNITIES,

Having regard to the Treaty establishing the European Community,

Having regard to Council Directive 89/107/EEC of 21 December 1988 on the approximation of the laws of the Member States concerning food additives authorized for use in foodstuffs intended for human consumption⁽¹⁾, as amended by European Parliament and Council Directive 94/34/EC⁽²⁾, and in particular Article 3 (3) (a) thereof,

After consulting the Scientific Committee for Food,

Whereas it is necessary to establish purity criteria for all additives other than colours and sweeteners mentioned in European Parliament and Council Directive 95/2/EC of 20 February 1995 on food additives other than colours and sweeteners⁽³⁾;

Whereas it is necessary to replace the purity criteria set out in Council Directive 65/66/EEC of 26 January 1965 laying down specific criteria of purity for preservatives authorized for use in foodstuffs intended for human consumption⁽⁴⁾, as last amended by Directive 86/604/EEC⁽⁵⁾;

Whereas it is necessary to replace the purity criteria set out in Council Directive 78/664/EEC of 25 July 1978 laying down specific criteria of purity for antioxidants which may be used in foodstuffs intended for

human consumption⁽⁶⁾, as amended by Directive 82/712/EEC⁽⁷⁾;

Whereas Directives 65/66/EEC and 78/664/EEC should be repealed accordingly;

Whereas it is necessary to take into account the specifications and analytical techniques for additives as set out in the *Codex Alimentarius* as drafted by the Joint FAO/WHO Expert Committee on Food Additives (Jecfa);

Whereas food additives, if prepared by production methods or starting materials significantly different from those included in the evaluation of the Scientific Committee for Food, or if different from those mentioned in this Directive, should be submitted for evaluation by the Scientific Committee for Food for the purposes of a full evaluation with emphasis on the purity criteria;

Whereas, the measures provided for in this Directive are in accordance with the opinion of the Standing Committee for Foodstuffs,

HAS ADOPTED THIS DIRECTIVE:

Article 1

The purity criteria referred to in Article 3 (3) (a) of Directive 89/107/EEC for food additives other than colours and sweeteners, as mentioned in Directive 95/2/EC, are set out in the Annex hereto.

Article 2

Directives 65/66/EEC and 78/664/EEC are hereby repealed.

⁽¹⁾ OJ No L 40, 11. 2. 1989, p. 27.

⁽²⁾ OJ No L 237, 10. 9. 1994, p. 1.

⁽³⁾ OJ No L 61, 18. 3. 1995, p. 1.

⁽⁴⁾ OJ No 22, 9. 2. 1965, p. 373.

⁽⁵⁾ OJ No L 352, 13. 12. 1986, p. 45.

⁽⁶⁾ OJ No L 223, 14. 8. 1978, p. 30.

⁽⁷⁾ OJ No L 297, 23. 10. 1982, p. 31.

Article 3

1. Member States shall bring into force the laws, regulations and administrative provisions necessary to comply with this Directive before 1 July 1997. They shall immediately inform the Commission thereof.

When Member States adopt these provisions, these shall contain a reference to this Directive or shall be accompanied by such reference at the time of their official publication. The procedure for such reference shall be adopted by Member States.

2. Products put on the market or labelled before 1 July 1997 which do not comply with this Directive may be marketed until stocks are exhausted.

Article 4

This Directive shall enter into force on the 20th day following that of its publication in the *Official Journal of the European Communities*.

Article 5

This Directive is addressed to the Member States.

Done at Brussels, 2 December 1996.

For the Commission

Martin BANGEMANN

Member of the Commission

ANNEX

E 200 SORBIC ACID

Definition

<i>Chemical name</i>	Sorbic acid Trans, trans-2,4-hexadienoic acid
Einecs	203-768-7
<i>Chemical formula</i>	C ₆ H ₈ O ₂
<i>Molecular weight</i>	112,12
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	Colourless needles or white free flowing powder, having a slight characteristic odour and showing no change in colour after heating for 90 minutes at 105 °C

Identification

A. Melting range	Between 133 °C and 135 °C, after vacuum drying for four hours in a sulphuric acid desiccator
B. Spectrometry	An isopropanol solution (1 in 4 000 000) shows absorbance maximum at 254 ±2 nm
C. Positive test for double bonds	
D. Sublimation point	80 °C

Purity

Water content	Not more than 0,5 % (Karl Fischer method)
Sulphated ash	Not more than 0,2 %
Aldehydes	Not more than 0,1 % (as formaldehyde)
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 202 POTASSIUM SORBATE

Definition

<i>Chemical name</i>	Potassium sorbate Potassium (E,E)-2,4-hexadienoate Potassium salt of trans, trans 2,4-hexadienoic acid
Einecs	246-376-1
<i>Chemical formula</i>	C ₆ H ₇ O ₂ K
<i>Molecular weight</i>	150,22
<i>Assay</i>	Content not less than 99 % on the dried basis
<i>Description</i>	White crystalline powder showing no change in colour after heating for 90 minutes at 105 °C

Identification

- A. Melting range of sorbic acid isolated by acidification and not re-crystallized 133°C to 135°C after vacuum drying in a sulphuric acid desiccator
- B. Positive tests for potassium and for double bonds

Purity

Loss on drying	Not more than 1,0 % (105°C, 3h)
Acidity or alkalinity	Not more than about 1,0 % (as sorbic acid or K ₂ CO ₃)
Aldehydes	Not more than 0,1 %, calculated as formaldehyde
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 203 CALCIUM SORBATE**Definition**

<i>Chemical name</i>	Calcium sorbate Calcium salts of trans, trans-2,4-hexadienoic acid
<i>Einecs</i>	231-321-6
<i>Chemical formula</i>	C ₁₂ H ₁₄ O ₄ Ca
<i>Molecular weight</i>	262,32
<i>Assay</i>	Content not less than 98 % on the dried basis
<i>Description</i>	Fine white crystalline powder not showing any change in colour after heating at 105°C for 90 minutes

Identification

- A. Melting range of sorbic acid isolated by acidification and not re-crystallized 133°C to 135°C after vacuum drying in a sulphuric acid desiccator
- B. Positive tests for calcium and for double bonds

Purity

Loss on drying	Not more than 2,0 %, determined by vacuum drying for four hours in a sulphuric acid desiccator
Aldehydes	Not more than 0,1 % (as formaldehyde)
Fluoride	Not more than 10 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 210 BENZOIC ACID

Definition

<i>Chemical name</i>	Benzoic acid Benzenecarboxylic acid Phenylcarboxylic acid
<i>Einecs</i>	200-618-2
<i>Chemical formula</i>	$C_7H_6O_2$
<i>Molecular weight</i>	122,12
<i>Assay</i>	Content not less than 99,5 % on the anhydrous basis
<i>Description</i>	White crystalline powder

Identification

A. Melting range	121,5 °C to 123,5 °C
B. Positive sublimation test and test for benzoate	

Purity

Loss on drying	Not more than 0,5 % after drying for three hours over sulphuric acid
pH	About 4 (solution in water)
Sulphated ash	Not more than 0,05 %
Chlorinated organic compounds	Not more than 0,07 % expressed as chloride corresponding to 0,3 % expressed as monochlorobenzoic acid
Readily oxidizable substances	Add 1,5 ml of sulphuric acid to 100 ml of water, heat to boiling point and add 0,1 N $KMnO_4$ in drops, until the pink colour persists for 30 seconds. Dissolve 1 g of the sample, weighed to the nearest mg, in the heated solution, and titrate with 0,1 N $KMnO_4$ to a pink colour that persists for 15 seconds. Not more than 0,5 ml should be required
Readily carbonizable substances	A cold solution of 0,5 g of benzoic acid in 5 ml of 94,5 to 95,5 % sulphuric acid must not show a stronger colouring than that of a reference liquid containing 0,2 ml of cobalt chloride TSC ⁽¹⁾ , 0,3 ml of ferric chloride TSC ⁽²⁾ , 0,1 ml of copper sulphate TSC ⁽³⁾ and 4,4 ml of water
Polycyclic acids	On fractional acidification of a neutralized solution of benzoic acid, the first precipitate must not have a different melting point from that of the benzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

⁽¹⁾ Cobalt chloride TSC: dissolve approximately 65 g of cobalt chloride $CoCl_2 \cdot 6H_2O$ in a sufficient quantity of a mixture of 25 ml hydrochloric acid and 975 ml of water to give a total volume of 1 litre. Place exactly 5 ml of this solution in a round-bottomed flask containing 250 ml of iodine solution, add 5 ml of 3 % hydrogen peroxide, then 15 ml of a 20 % solution of sodium hydroxide. Boil for 10 minutes, allow to cool, add 2 g of potassium iodide and 20 ml of 25 % sulphuric acid. After the precipitate is completely dissolved, titrate the liberated iodine with sodium thiosulphate (0,1 N) in the presence of starch TS(*). 1 ml of sodium thiosulphate (0,1 N) corresponds to 23,80 mg of $CoCl_2 \cdot 6H_2O$. Adjust final volume of solution by the addition of a sufficient quantity of the hydrochloric acid/water mixture to give a solution containing 59,5 mg of $CoCl_2 \cdot 6H_2O$ per ml.

⁽²⁾ Ferric chloride TSC: dissolve approximately 55 g of ferric chloride in a sufficient quantity of a mixture of 25 ml of hydrochloric acid and 975 ml of water to give a total volume of 1 litre. Place 10 ml of this solution in a round-bottomed flask containing 250 ml of iodine solution, add 15 ml of water and 3 g of potassium iodide; leave the mixture to stand for 15 minutes. Dilute with 100 ml of water then titrate the liberated iodine with sodium thiosulphate (0,1 N) in the presence of starch TS(*). 1 ml of sodium thiosulphate (0,1 N) corresponds to 27,03 mg of $FeCl_3 \cdot 6H_2O$. Adjust final volume of solution by the addition of a sufficient quantity of the hydrochloric acid/water to give a solution containing 45,0 mg of $FeCl_3 \cdot 6H_2O$ per ml.

⁽³⁾ Copper sulphate TSC: dissolve approximately 65 g of copper sulphate $CuSO_4 \cdot 5H_2O$ in a sufficient quantity of a mixture of 25 ml of hydrochloric acid and 975 ml of water to give a total volume of 1 litre. Place 10 ml of this solution in a round-bottomed flask containing 250 ml of iodine solution, add 40 ml of water, 4 ml of acetic acid and 3 g of potassium iodide. Titrate the liberated iodine with sodium thiosulphate (0,1 N) in the presence of starch TS(*). 1 ml of sodium thiosulphate (0,1 N) corresponds to 24,97 mg of $CuSO_4 \cdot 5H_2O$. Adjust final volume of solution by the addition of a sufficient quantity of the hydrochloric acid/water mixture to give a solution containing 62,4 mg of $CuSO_4 \cdot 5H_2O$ per ml.

(*) Starch TS: triturate 0,5 g starch (potato starch, maize starch of soluble starch) with 5 ml of water; to the resulting paste add a sufficient quantity of water to give a total volume of 100 ml, stirring all the time. Boil for a few minutes, allow to cool, filter. The starch must be freshly prepared.

E 211 SODIUM BENZOATE

Definition

Chemical name

Sodium benzoate
Sodium salt of benzenecarboxylic acid
Sodium salt of phenylcarboxylic acid

Einecs

208-534-8

Chemical formula $C_7H_5O_2Na$ *Molecular weight*

144,11

*Assay*Not less than 99 % of $C_7H_5O_2Na$, after drying at 105 °C for four hours*Description*

A white, almost odourless, crystalline powder or granules

Identification

A. Solubility

Freely soluble in water, sparingly soluble in ethanol

B. Melting range for benzoic acid

Melting range of benzoic acid isolated by acidification and not recrystallized 121,5 °C to 123,5 °C, after drying in a sulphuric acid desiccator

C. Positive tests for benzoate and for sodium

Purity

Loss on drying

Not more than 1,5 % after drying at 105 °C for four hours

Readily oxidizable substances

Add 1,5 ml of sulphuric acid to 100 ml of water, heat to boiling point and add 0,1 N $KMnO_4$ in drops, until the pink colour persists for 30 seconds. Dissolve 1 g of the sample, weighed to the nearest mg, in the heated solution, and titrate with 0,1 N $KMnO_4$ to a pink colour that persists for 15 seconds. Not more than 0,5 ml should be required

Polycyclic acids

On fractional acidification of a (neutralized) solution of sodium benzoate, the first precipitate must not have a different melting range from that of benzoic acid

Chlorinated organic compounds

Not more than 0,06 % expressed as chloride, corresponding to 0,25 % expressed as monochlorobenzoic acid

Degree of acidity or alkalinity

Neutralization of 1 g of sodium benzoate, in the presence of phenolphthalein, must not require more than 0,25 ml of 0,1 N NaOH or 0,1 N HCl

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 212 POTASSIUM BENZOATE

Definition

Chemical name

Potassium benzoate
Potassium salt of benzenecarboxylic acid
Potassium salt of phenylcarboxylic acid

Einecs

209-481-3

Chemical formula $C_7H_5KO_2 \cdot 3H_2O$

<i>Molecular weight</i>	214,27
<i>Assay</i>	Content not less than 99 % $C_7H_5O_2K$ after drying at 105 °C to constant weight
<i>Description</i>	White crystalline powder
Identification	
A. Melting range of benzoic acid isolated by acidification and not recrystallized 121,5 °C to 123,5 °C, after vacuum drying in a sulphuric acid desiccator	
B. Positive tests for benzoate and for potassium	
Purity	
Loss on drying	Not more than 26,5 %, determined by drying at 105 °C
Chlorinated organic compounds	Not more than 0,06 % expressed as chloride, corresponding to 0,25 % expressed as monochlorobenzoic acid
Readily oxidizable substances	Add 1,5 ml of sulphuric acid to 100 ml of water, heat to boiling point and add 0,1 N $KMnO_4$ in drops, until the pink colour persists for 30 seconds. Dissolve 1 g of the sample, weighed to the nearest mg, in the heated solution, and titrate with 0,1 N $KMnO_4$ to a pink colour that persists for 15 seconds. Not more than 0,5 ml should be required
Readily carbonizable substances	A cold solution of 0,5 g of benzoic acid in 5 ml 94,5 to 95,5 % sulphuric acid must not show a stronger colouring than that of a reference liquid containing 0,2 ml of cobalt chloride TSC, 0,3 ml of ferric chloride TSC, 0,1 ml of copper sulphate TSC and 4,4 ml of water
Polycyclic acids	On fractional acidification of a (neutralized) solution of potassium benzoate, the first precipitate must not have a different melting range from that of benzoic acid
Degree of acidity or alkalinity	Neutralization of 1 g of potassium benzoate, in the presence of phenolphthalein, must not require more than 0,25 ml of 0,1 N NaOH or 0,1 N HCl
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 213 CALCIUM BENZOATE

Synonyms	Monocalcium benzoate
Definition	
<i>Chemical name</i>	Calcium benzoate Calcium dibenzoate

Einecs	218-235-4
<i>Chemical formula</i>	Anhydrous: $C_{14}H_{10}O_4Ca$ Monohydrate: $C_{14}H_{10}O_4Ca \cdot H_2O$ Trihydrate: $C_{14}H_{10}O_4Ca \cdot 3H_2O$
<i>Molecular weight</i>	Anhydrous: 282,31 Monohydrate: 300,32 Trihydrate: 336,36
<i>Assay</i>	Content not less than 99 % after drying at 105 °C
<i>Description</i>	White or colourless crystals, or white powder
Identification	
A. Melting range of benzoic acid isolated by acidification and not recrystallized 121,5 °C to 123,5 °C, after vacuum drying in a sulphuric acid desiccator	
B. Positive tests for benzoate and for calcium	
Purity	
Loss on drying	Not more than 17,5 % determined by drying at 105 °C to constant weight
Water insoluble matter	Not more than 0,3 %
Chlorinated organic compounds	Not more than 0,06 % expressed as chloride, corresponding to 0,25 % expressed as monochlorobenzoic acids
Readily oxidizable substances	Add 1,5 ml of sulphuric acid to 100 ml of water, heat to boiling point and add 0,1 N $KMnO_4$ in drops, until the pink colour persists for 30 seconds. Dissolve 1 g of the sample, weighed to the nearest mg, in the heated solution, and titrate with 0,1 N $KMnO_4$ to a pink colour that persists for 15 seconds. Not more than 0,5 ml should be required
Readily carbonizable substances	Cold solution of 0,5 g of benzoic acid in 5 ml of 94,5 to 95,5 % sulphuric acid must not show a stronger colouring than that of a reference liquid containing 0,2 ml of cobalt chloride TSC, 0,3 ml of ferric chloride TSC, 0,1 ml of copper sulphate TSC and 4,4 ml of water
Polycyclic acids	On fractional acidification of a (neutralized) solution of calcium benzoate, the first precipitate must not be a different melting range from that of benzoic acid
Degree of acidity or alkalinity	Neutralization of 1 g of calcium benzoate, in the presence of phenolphthalein, must not require more than 0,25 ml of 0,1 N NaOH or 0,1 N HCl
Fluoride	Not more than 10 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 214 ETHYL *p*-HYDROXYBENZOATE

Synonyms	Ethylparaben Ethyl <i>p</i> -oxybenzoate
-----------------	---

Definition

<i>Chemical name</i>	Ethyl- <i>p</i> -hydroxybenzoate Ethyl ester of <i>p</i> -hydroxybenzoic acid
Einecs	204-399-4
<i>Chemical formula</i>	C ₉ H ₁₀ O ₃
<i>Molecular weight</i>	166,8
<i>Assay</i>	Content not less than 99,5 % after drying for two hours at 80 °C
<i>Description</i>	Almost odourless, small, colourless crystals or a white, crystalline powder

Identification

A. Melting range	115 °C to 118 °C
B. Positive test for <i>p</i> -hydroxybenzoate	Melting range of <i>p</i> -hydroxybenzoic acid isolated by acidification and not recrystallized: 213 °C to 217 °C, after vacuum drying in a sulphuric acid desiccator
C. Positive test for alcohol	

Purity

Loss on drying	Not more than 0,5 % after drying for two hours at 80 °C
Sulphated ash	Not more than 0,05 %
<i>p</i> -Hydroxybenzoic acid and salicylic acid	Not more than 0,35 % expressed as <i>p</i> -hydroxybenzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 215 SODIUM ETHYL *p*-HYDROXYBENZOATE**Definition**

<i>Chemical name</i>	Sodium ethyl <i>p</i> -hydroxybenzoate Sodium compound of the ethyl ester of <i>p</i> -hydroxybenzoic acid
Einecs	252-487-6
<i>Chemical formula</i>	C ₉ H ₉ O ₃ Na
<i>Molecular weight</i>	188,8
<i>Assay</i>	Content of ethylester of <i>p</i> -hydroxybenzoic acid not less than 83 % on the anhydrous basis
<i>Description</i>	White, crystalline hygroscopic powder

Identification

A. Melting range	115 °C to 118 °C, after vacuum drying in a sulphuric acid desiccator
------------------	--

B. Positive test for <i>p</i> -hydroxybenzoate	Melting range of <i>p</i> -hydroxybenzoic acid derived from the sample is 213°C to 217°C
C. Positive test for sodium	
D. pH of a 0,1% aqueous solution must be between 9,9 and 10,3	
Purity	
Loss on drying	Not more than 5 %, determined by vacuum drying in a sulphuric acid desiccator
Sulphated ash	37 to 39 %
<i>p</i> -Hydroxybenzoic acid and salicylic acid	Not more than 0,35 % expressed as <i>p</i> -hydroxybenzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 216 PROPYL *p*-HYDROXYBENZOATE

Synonyms	Propylparaben Propyl <i>p</i> -oxybenzoate
Definition	
<i>Chemical name</i>	Propyl <i>p</i> -hydroxybenzoate n-Propyl <i>p</i> -hydroxybenzoic acid
Einecs	202-307-7
<i>Chemical formula</i>	C ₁₀ H ₁₂ O ₃
<i>Molecular weight</i>	180,21
<i>Assay</i>	Content not less than 99,5 % after drying for two hours at 80°C
<i>Description</i>	Almost odourless, small, colourless crystals or a white, crystalline powder
Identification	
A. Melting range	95°C to 97°C after drying for two hours at 80°C
B. Positive test for <i>p</i> -hydroxybenzoate	Melting range of <i>p</i> -hydroxybenzoic acid derived from the sample is 213°C to 217°C
Purity	
Loss on drying	Not more than 0,5 % after drying for two hours at 80°C
Sulphated ash	Not more than 0,05 %
<i>p</i> -Hydroxybenzoic acid and salicylic acid	Not more than 0,35 % expressed as <i>p</i> -hydroxybenzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 217 SODIUM PROPYL *p*-HYDROXYBENZOATE

Definition

Chemical name	Sodium n-propyl <i>p</i> -hydroxybenzoate Sodium compound of the n-propylester of <i>p</i> -hydroxybenzoic acid
Einecs	252-488-1
Chemical formula	C ₁₀ H ₁₁ O ₃ Na
Molecular weight	202,21
Assay	Content of the propyl ester of <i>p</i> -hydroxybenzoic acid not less than 85 % on the anhydrous basis
Description	White, or almost white, crystalline hygroscopic powder

Identification

- A. Melting range of ester isolated by acidification and not recrystallized: 94°C to 97°C, after vacuum drying in a sulphuric acid desiccator
- B. Positive test for sodium
- C. pH of a 0,1 % aqueous solution must be between 9,8 and 10,2

Purity

Loss on drying	Not more than 5 %, determined by vacuum drying in a sulphuric acid desiccator
Sulphated ash	34 to 36 %
<i>p</i> -Hydroxybenzoic acid and salicylic acid	Not more than 0,35 % expressed as <i>p</i> -hydroxybenzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 218 METHYL *p*-HYDROXYBENZOATE

Synonyms

Methylparaben
Methyl-*p*-oxybenzoate

Definition

Chemical name	Methyl <i>p</i> -hydroxybenzoate Methyl ester of <i>p</i> -hydroxybenzoic acid
Einecs	243-171-5
Chemical formula	C ₈ H ₈ O ₃

<i>Molecular weight</i>	152,15
<i>Assay</i>	Content not less than 99 % after drying for two hours at 80 °C
<i>Description</i>	Almost odourless, small colourless crystals or white crystalline powder
Identification	
A. Melting range	125 °C to 128 °C
B. Positive test for <i>p</i> -hydroxybenzoate	Melting range of <i>p</i> -hydroxybenzoic acid derived from the sample is 213 °C to 217 °C after drying for two hours at 80 °C
Purity	
Loss on drying	Not more than 0,5 %, after drying for two hours at 80 °C
Sulphated ash	Not more than 0,05 %
<i>p</i> -Hydroxybenzoic acid and salicylic acid	Not more than 0,35 % expressed as <i>p</i> -hydroxybenzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 219 SODIUM METHYL *p*-HYDROXYBENZOATE

Definition	
<i>Chemical name</i>	Sodium methyl <i>p</i> -hydroxybenzoate Sodium compound of the methylester of <i>p</i> -hydroxybenzoic acid
<i>Chemical formula</i>	C ₈ H ₇ O ₃ Na
<i>Molecular weight</i>	174,15
<i>Assay</i>	Content not less than 99,5 % on the anhydrous basis
<i>Description</i>	White, hygroscopic powder
Identification	
A. The white precipitate formed by acidifying with hydrochloric acid a 10 % (w/v) aqueous solution of the sodium derivative of methyl <i>p</i> -hydroxybenzoate (using litmus paper as indicator) shall, when washed with water and dried at 80 °C for two hours, have a melting range of 125 °C to 128 °C	
B. Positive test for sodium	
C. pH of a 0,1 % solution in carbon dioxide free water, not less than 9,7 and not more than 10,3	

Purity

Water content	Not more than 5 % (Karl Fischer method)
Sulphated ash	40 % to 44,5 % on the anhydrous basis
<i>p</i> -Hydroxybenzoic acid and salicylic acid	Not more than 0,35 % expressed as <i>p</i> -hydroxybenzoic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 220 SULPHUR DIOXIDE**Definition**

<i>Chemical name</i>	Sulphur dioxide Sulphurous acid anhydride
<i>Einecs</i>	231-195-2
<i>Chemical formula</i>	SO ₂
<i>Molecular weight</i>	64,07
<i>Assay</i>	Content not less than 99 %
<i>Description</i>	Colourless, non-flammable gas with strong pungent suffocating odour

Identification

A. Positive test for sulphurous substances

Purity

Water content	Not more than 0,05 %
Non-volatile residue	Not more than 0,01 %
Sulphur trioxide	Not more than 0,1 %
Selenium	Not more than 10 mg/kg
Other gases not normally present in the air	No trace
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 221 SODIUM SULPHITE

Definition

<i>Chemical name</i>	Sodium sulphite (anhydrous or heptahydrate)
<i>Einecs</i>	231-821-4
<i>Chemical formula</i>	Anhydrous: Na_2SO_3 Heptahydrate: $\text{Na}_2\text{SO}_3 \cdot 7\text{H}_2\text{O}$
<i>Molecular weight</i>	Anhydrous: 126,04 Heptahydrate: 252,16
<i>Assay</i>	Anhydrous: Not less than 95 % of Na_2SO_3 and not less than 48 % of SO_2 Heptahydrate: Not less than 48 % of Na_2SO_3 and not less than 24 % of SO_2
<i>Description</i>	White crystalline powder or colourless crystals

Identification

- A. Positive tests for sulphite and for sodium
- B. pH of a 10 % solution (anhydrous) or a 20 % solution (heptahydrate) between 8,5 and 11,5

Purity

Thiosulphate	Not more than 0,1 % based on the SO_2 content
Iron	Not more than 50 mg/kg based on the SO_2 content
Selenium	Not more than 10 mg/kg based on the SO_2 content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 222 SODIUM BISULPHITE

Definition

<i>Chemical name</i>	Sodium bisulphite Sodium hydrogen sulphite
<i>Einecs</i>	231-921-4
<i>Chemical formula</i>	NaHSO_3 in aqueous solution
<i>Molecular weight</i>	104,06
<i>Assay</i>	Content not less than 32 % w/w NaHSO_3
<i>Description</i>	A clear, colourless to yellow solution

Identification

- A. Positive tests for sulphite and for sodium
- B. pH of a 10 % aqueous solution between 2,5 and 5,5

Purity

Iron	Not more than 50 mg/kg of Na ₂ SO ₃ based on the SO ₂ content
Selenium	Not more than 10 mg/kg based on the SO ₂ content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 223 SODIUM METABISULPHITE**Synonyms**

Pyrosulphite
Sodium pyrosulphite

Definition

<i>Chemical name</i>	Sodium disulphite Disodium pentaoxodisulphate
<i>Einecs</i>	231-673-0
<i>Chemical formula</i>	Na ₂ S ₂ O ₅
<i>Molecular weight</i>	190,11
<i>Assay</i>	Content not less than 95 % Na ₂ S ₂ O ₅ and not less than 64 % of SO ₂
<i>Description</i>	White crystals or crystalline powder

Identification

- A. Positive tests for sulphite and for sodium
- B. pH of a 10 % aqueous solution between 4,0 and 5,5

Purity

Thiosulphate	Not more than 0,1 % based on the SO ₂ content
Iron	Not more than 50 mg/kg based on the SO ₂ content
Selenium	Not more than 10 mg/kg based on the SO ₂ content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg

Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 224 POTASSIUM METABISULPHITE

Synonyms	Potassium pyrosulphite
Definition	
<i>Chemical name</i>	Potassium disulphite Potassium pentaoxo disulphate
Einecs	240-795-3
<i>Chemical formula</i>	$K_2S_2O_5$
<i>Molecular weight</i>	222,33
<i>Assay</i>	Content not less than 90 % of $K_2S_2O_5$ and not less than 51,8 % of SO_2 , the remainder being composed almost entirely of potassium sulphate
<i>Description</i>	Colourless crystals or white crystalline powder
Identification	
A. Positive tests for sulphite and for potassium	
Purity	
Thiosulphate	Not more than 0,1 % based on the SO_2 content
Iron	Not more than 50 mg/kg based on the SO_2 content
Selenium	Not more than 10 mg/kg based on the SO_2 content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 226 CALCIUM SULPHITE

Definition	
<i>Chemical name</i>	Calcium sulphite
Einecs	218-235-4
<i>Chemical formula</i>	$CaSO_3 \cdot 2H_2O$
<i>Molecular weight</i>	156,17
<i>Assay</i>	Content not less than 95 % of $CaSO_3 \cdot 2H_2O$ and not less than 39 % of SO_2
<i>Description</i>	White crystals or white crystalline powder

Identification

- A. Positive tests for sulphite and for calcium

Purity

Iron	Not more than 50 mg/kg based on the SO ₂ content
Selenium	Not more than 10 mg/kg based on the SO ₂ content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 227 CALCIUM BISULPHITE**Definition**

<i>Chemical name</i>	Calcium bisulphite Calcium hydrogen sulphite
Einecs	237-423-7
<i>Chemical formula</i>	Ca(HSO ₃) ₂
<i>Molecular weight</i>	202,22
<i>Assay</i>	6 to 8 % (w/v) of sulphur dioxide and 2,5 to 3,5 % (w/v) of calcium dioxide corresponding to 10 to 14 % (w/v) of calcium bisulphite [Ca(HSO ₃) ₂]
<i>Description</i>	Clear greenish-yellow aqueous solution having a distinct odour of sulphur dioxide

Identification

- A. Positive tests for sulphite and for calcium

Purity

Iron	Not more than 50 mg/kg based on the SO ₂ content
Selenium	Not more than 10 mg/kg based on the SO ₂ content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 228 POTASSIUM BISULPHITE**Definition**

<i>Chemical name</i>	Potassium bisulphite Potassium hydrogen sulphite
----------------------	---

Einecs	231-870-1
<i>Chemical formula</i>	KHSO ₃ in aqueous solution
<i>Molecular weight</i>	120,17
<i>Assay</i>	Content not less than 280 g KHSO ₃ per litre (or 150 g SO ₂ per litre)
<i>Description</i>	Clear colourless aqueous solution

Identification

- A. Positive tests for sulphite and for potassium

Purity

Iron	Not more than 50 mg/kg based on the SO ₂ content
Selenium	Not more than 10 mg/kg based on the SO ₂ content
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 230 BIPHENYL**Synonyms**

Diphenyl

Definition

Chemical name
1,1'-biphenyl
Phenylbenzene

Einecs
202-163-5

Chemical formula
C₁₂H₁₀

Molecular weight
154,20

Assay
Content not less than 99,8 %

Description
White or pale yellow to amber crystalline solid having a characteristic odour

Identification

- A. Melting range
68,5°C to 70,5°C
- B. Distillation range
It distils completely within a 2,5°C range between 252,5°C and 257,5°C

Purity

Benzene	Not more than 10 mg/kg
Aromatic amines	Not more than 2 mg/kg (as aniline)
Phenol derivatives	Not more than 5 mg/kg (as phenol)

Readily carbonizable substances	Cold solution of 0,5 g of biphenyl in 5 ml of 94,5 to 95,5 % sulphuric acid must not show a stronger colouring than that of a reference liquid containing 0,2 ml of cobalt chloride TSC, 0,3 ml of ferric chloride TSC, 0,1 ml of copper sulphate TSC and 4,4 ml of water
Terphenyl and higher polyphenyl derivatives	Not more than 0,2 %
Polycyclic aromatic hydrocarbons	Absent
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 231 ORTHOPHENYLPHENOL

Synonyms	Orthoxenol
Definition	
<i>Chemical name</i>	(1,1'-Biphenyl)-2-ol 2-Hydroxydiphenyl o-Hydroxydiphenyl
Einecs	201-993-5
<i>Chemical formula</i>	C ₁₂ H ₁₀ O
<i>Molecular weight</i>	170,20
<i>Assay</i>	Content not less than 99 %
<i>Description</i>	White or slightly yellowish crystalline powder
Identification	
A. Melting range	56 °C to 58 °C
B. Positive test for phenolate	An ethanolic solution (1 g in 10 ml) produces a green colour on addition of 10 % ferric chloride solution
Purity	
Sulphated ash	Not more than 0,05 %
Diphenyl ether	Not more than 0,3 %
<i>p</i> -Phenylphenol	Not more than 0,1 %
1-Naphthol	Not more than 0,01 %
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 232 SODIUM ORTHOPHENYLPHENOL

Synonyms

Sodium orthophenylphenate
Sodium salt of o-phenylphenol

Definition

Chemical name

Sodium orthophenylphenol

Einecs

205-055-6

Chemical formula $C_{12}H_9ONa \cdot 4H_2O$ *Molecular weight*

264,26

*Assay*Content not less than 97 % of $C_{12}H_9ONa \cdot 4H_2O$ *Description*

White or slightly yellowish crystalline powder

Identification

- A. Positive tests for phenolate and for sodium
- B. Melting range of orthophenylphenol isolated by acidification and not recrystallized derived from the sample 56 °C to 58 °C after drying in a sulphuric acid desiccator
- C. pH of a 2 % aqueous solution must be between 11,1 and 11,8

Purity

Diphenylether

Not more than 0,3 %

p-phenylphenol

Not more than 0,1 %

1-naphthol

Not more than 0,01 %

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 233 THIABENDAZOLE

Definition

Chemical name

4-(2-benzimidazolyl)thiazole
2-(4-thiazolyl)-1H-benzimidazole

Einecs

1205-725-8

Chemical formula $C_{10}H_7N_3S$

<i>Molecular weight</i>	201,26
<i>Assay</i>	Content not less than 98 % on the anhydrous basis
<i>Description</i>	White, or almost white, odourless powder
Identification	
A. Melting range	296 °C to 303 °C
B. Spectrometry	Absorption maxima in 0,1 N HCl (0,0005 % w/v) at 302 nm, 258 nm and 243 nm $E_{1\text{ cm}}^{1\%}$ at 302 nm ± 2 nm: approximately 1 230 $E_{1\text{ cm}}^{1\%}$ at 258 nm ± 2 nm: approximately 200 $E_{1\text{ cm}}^{1\%}$ at 243 nm ± 2 nm: approximately 620 Ratio of absorption 243 nm/302 nm = 0,47 to 0,53 Ratio of absorption 258 nm/302 nm = 0,14 to 0,18
Purity	
Water content	Not more than 0,5 % (Karl Fischer method)
Sulphated ash	Not more than 0,2 %
Selenium	Not more than 3 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 234 NISIN

Definition	Nisin consists of several closely related polypeptides produced by natural strains of <i>Streptococcus lactis</i> , Lancefield group N
Einecs	215-807-5
<i>Chemical formula</i>	$C_{143}H_{230}N_{42}O_{37}S_7$
<i>Molecular weight</i>	3 354,12
<i>Assay</i>	Nisin concentrate contains not less than 900 units per mg in a mixture of non-fat milk solids and a minimum sodium chloride content of 50 %
<i>Description</i>	White powder
Purity	
Loss on drying	Not more than 3 % when dried to constant weight at 102 °C to 103 °C
Arsenic	Not more than 1 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 235 NATAMYCIN

Synonyms

Pimaricin

Definition

Natamycin is a fungicide of the polyene macrolide group, and is produced by natural strains of *Streptomyces natalensis* or of *Streptococcus lactis*

Einecs

231-683-5

Chemical formula

 $C_{33}H_{47}O_{13}N$

Molecular weight

665,74

Assay

Content not less than 95 % on the anhydrous basis

Description

White to creamy-white crystalline powder

Identification

A. Colour reactions

On adding a few crystals of natamycin on a spot plate, to a drop of:

— concentrated hydrochloric acid, a blue colour develops,

— concentrated phosphoric acid, a green colour develops,

which changes into pale red after a few minutes

B. Spectrometry

A 0,0005 % w/v solution in 1 % methanolic acetic acid solution has absorption maxima at about 290 nm, 303 nm and 318 nm, a shoulder at about 280 nm and exhibits minima at about 250 nm, 295,5 nm and 311 nm

C. pH

5,5 to 7,5 (1 % w/v solution in previously neutralized mixture of 20 parts dimethylformamide and 80 parts of water)

D. Specific rotation

$[\alpha]_D^{20} = +250^\circ$ to $+295^\circ$ (a 1 % w/v solution in glacial acetic acid, at 20°C and calculated with reference to the dried material)

Purity

Loss on drying

Not more than 8 % (over P_2O_5 , in vacuum at 60°C to constant weight)

Sulphated ash

Not more than 0,5 %

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

Microbiological criteria: total viable count

Not more than 100/g

E 239 HEXAMETHYLENE TETRAMINE

Synonyms

Hexamine
Methenamine

Definition

Chemical name

1,3,5,7-Tetraazatricyclo [3.3.1.1^{3,7}]-decane, hexamethylenetetramine

Einecs

202-905-8

<i>Chemical formula</i>	$C_6H_{12}N_4$
<i>Molecular weight</i>	140,19
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	Colourless or white crystalline powder
Identification	
A. Positive tests for formaldehyde and for ammonia	
B. Sublimation point approximately 260 °C	
Purity	
Loss on drying	Not more than 0,5 % after drying at 105 °C in vacuum over P_2O_5 for two hours
Sulphated ash	Not more than 0,05 %
Sulphates	Not more than 0,005 % expressed as SO_4
Chlorides	Not more than 0,005 % expressed as Cl
Ammonium salts	Not detectable
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 242 DIMETHYL DICARBONATE

Synonyms	DMDC Dimethyl pyrocarbonate
Definition	
Einecs	224-859-8
<i>Chemical name</i>	Dimethyl dicarbonate Pyrocarbonic acid dimethyl ester
<i>Chemical formula</i>	$C_4H_6O_5$
<i>Molecular weight</i>	134,09
<i>Assay</i>	Content not less than 99,8 %
<i>Description</i>	Colourless liquid, decomposes in aqueous solution. It is corrosive to skin and eyes and toxic by inhalation and ingestion

Identification

A. Decomposition	After dilution positive tests for CO ₂ and methanol
B. Melting point	17°C
Boiling point	172°C with decomposition
C. Density 20°C	Approximately 1,25 g/cm ³
D. Infrared spectrum	Maxima at 1 156 and 1 832 cm ⁻¹

Purity

Dimethyl carbonate	Not more than 0,2 %
Chlorine, total	Not more than 3 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 249 POTASSIUM NITRITE**Definition**

<i>Chemical name</i>	Potassium nitrite
Einecs	231-832-4
<i>Chemical formula</i>	KNO ₂
<i>Molecular weight</i>	85,11
<i>Assay</i>	Content not less than 95 % on the anhydrous basis ⁽¹⁾
<i>Description</i>	White or slightly yellow, deliquescent granules

Identification

- A. Positive tests for nitrite and for potassium
- B. pH of a 5 % solution: not less than 6,0 and not more than 9,0

Purity

Loss on drying	Not more than 3 % after drying for four hours over silica gel
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

⁽¹⁾ When labelled 'for food use', nitrite may only be sold in a mixture with salt or a salt substitute.

E 250 SODIUM NITRITE

Definition

<i>Chemical name</i>	Sodium nitrite
Einecs	231-555-9
<i>Chemical formula</i>	NaNO_2
<i>Molecular weight</i>	69,00
<i>Assay</i>	Content not less than 97 % on the anhydrous basis ⁽¹⁾
<i>Description</i>	White crystalline powder or yellowish lumps

Identification

- A. Positive tests for nitrite and for sodium

Purity

Loss on drying	Not more than 0,25 % after drying over silica gel for four hours
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 251 SODIUM NITRATE

Synonyms

Chile saltpetre
Cubic or soda nitre

Definition

<i>Chemical name</i>	Sodium nitrate
Einecs	231-554-3
<i>Chemical formula</i>	NaNO_3
<i>Molecular weight</i>	85,00
<i>Assay</i>	Content not less than 99 % after drying at 105 °C for four hours
<i>Description</i>	White crystalline, slightly hygroscopic powder

Identification

- A. Positive tests for nitrate and for sodium
- B. pH of a 5 % solution
- C. Melting point: $\pm 308^\circ\text{C}$
- Not less than 5,5 and more than 8,3

⁽¹⁾ When labelled 'for food use', nitrite may only be sold in a mixture with salt or a salt substitute.

Purity

Loss on drying	Not more than 2 % after drying at 105 °C for four hours
Nitrites	Not more than 30 mg/kg expressed as NaNO ₂
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 252 POTASSIUM NITRATE**Synonyms**

Chile saltpetre
Cubic or soda nitre

Definition

<i>Chemical name</i>	Potassium nitrate
<i>Einecs</i>	231-818-8
<i>Chemical formula</i>	KNO ₃
<i>Molecular weight</i>	101,11
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	White crystalline powder or transparent prisms having a cooling, saline, pungent taste

Identification

- | | |
|---|---|
| A. Positive tests for nitrate and for potassium | |
| B. pH of a 5 % solution | Not less than 4,5 and not more than 8,5 |

Purity

Loss on drying	Not more than 1 % after drying at 105 °C for four hours
Nitrites	Not more than 20 mg/kg expressed as KNO ₂
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 260 ACETIC ACID**Definition**

<i>Chemical name</i>	Acetic acid Ethanoic acid
----------------------	------------------------------

Einecs	200-580-7
<i>Chemical formula</i>	$C_2H_4O_2$
<i>Molecular weight</i>	60,05
<i>Assay</i>	Content not less than 99,8 %
<i>Description</i>	Clear, colourless liquid having a pungent, characteristic odour
Identification	
A. Boiling point	118°C at 760 mm pressure (of mercury)
B. Specific gravity	About 1,049
C. A one in three solution gives positive tests for acetate	
D. Solidification point	Not lower than 14,5°C
Purity	
Non-volatile residue	Not more than 100 mg/kg
Formic acid, formates and other oxidizable substances	Not more than 1 000 mg/kg expressed as formic acid
Readily oxidizable substances	Dilute 2 ml of the sample in a glass-stoppered container with 10 ml of water and add 0,1 ml of 0,1 N potassium permanganate. The pink colour does not change to brown within 30 minutes
Arsenic	Not more than 1 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 261 POTASSIUM ACETATE

Definition	
<i>Chemical name</i>	Potassium acetate
Einecs	204-822-2
<i>Chemical formula</i>	$C_2H_3O_2K$
<i>Molecular weight</i>	98,14
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	Colourless, deliquescent crystals or a white crystalline powder, odourless or with a faint acetic odour
Identification	
A. pH of a 5 % aqueous solution	Not less than 7,5 and not more than 9,0
B. Positive tests for acetate and for potassium	

Purity

Loss on drying	Not more than 8 % after drying at 150°C for two hours
Formic acid, formates and other oxidizable substances	Not more than 1 000 mg/kg expressed as formic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 262 (i) SODIUM ACETATE**Definition**

<i>Chemical name</i>	Sodium acetate
<i>Einecs</i>	204-823-8
<i>Chemical formula</i>	$C_2H_3NaO_2 \cdot nH_2O$ (n = 0 or 3)
<i>Molecular weight</i>	Anhydrous: 82,03 Trihydrate: 136,08
<i>Assay</i>	Content (for both of anhydrous and trihydrate form) not less than 98,5 % on the anhydrous basis
<i>Description</i>	Anhydrous: White, odourless, granular, hygroscopic powder Trihydrate: Colourless, transparent crystals or a granular crystalline powder, odourless or with a faint, acetic odour. Effloresces in warm, dry air

Identification

A. pH of a 1 % aqueous solution	Not less than 8,0 and not more than 9,5
B. Positive tests for acetate and for sodium	

Purity

Loss on drying	Anhydrous: Not more than 2 % (120°C, 4 hours) Trihydrate: Between 36 and 42 % (120°C, 4 hours)
Formic acid, formates and other oxidizable substances	Not more than 1 000 mg/kg expressed as formic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 262 (ii) SODIUM DIACETATE

Definition

Sodium diacetate is a molecular compound of sodium acetate and acetic acid

Chemical name

Sodium hydrogen diacetate

Einecs

204-814-9

Chemical formula

$C_4H_7NaO_4 \cdot nH_2O$ ($n = 0$ or 3)

Molecular weight

142,09 (anhydrous)

Assay

Content 39 to 41 % of free acetic acid and 58 to 60 % of sodium acetate

Description

White, hygroscopic crystalline solid with an acetic odour

Identification

A. pH of a 10 % aqueous solution

Not less than 4,5 and not more than 5,0

B. Positive tests for acetate and for sodium

Purity

Water content

Not more than 2 % (Karl Fischer method)

Formic acid, formates and other oxidizable substances

Not more than 1 000 mg/kg expressed as formic acid

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 263 CALCIUM ACETATE

Definition

Chemical name

Calcium acetate

Einecs

200-540-9

Chemical formula

Anhydrous: $C_4H_6O_4Ca$
Monohydrate: $C_4H_6O_4Ca \cdot H_2O$

Molecular weight

Anhydrous: 158,17
Monohydrate: 176,18

Assay

Content not less than 98 % on the anhydrous basis

Description

Anhydrous calcium acetate is a white, hygroscopic, bulky, crystalline solid with a slightly bitter taste. A slight odour of acetic acid may be present. The monohydrate may be needles, granules or powder

Identification

A. pH of a 10 % aqueous solution

Not less than 6,0 and not more than 9,0

B. Positive tests for acetate and for calcium

Purity

Loss on drying	Not more than 11 % after drying (155 °C to constant weight, for the monohydrate)
Water insoluble matter	Not more than 0,3 %
Formic acid, formates and other oxidizable substances	Not more than 1 000 mg/kg expressed as formic acid
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 270 LACTIC ACID**Definition**

<i>Chemical name</i>	Lactic acid 2-Hydroxypropionic acid 1-Hydroxyethane-1-carboxylic acid
<i>Einecs</i>	200-018-0
<i>Chemical formula</i>	$C_3H_6O_3$
<i>Molecular weight</i>	90,08
<i>Assay</i>	Content not less than 76 % and not more than 84 %
<i>Description</i>	Colourless or yellowish, nearly odourless, syrupy liquid with an acid taste, consisting of a mixture of lactic acid ($C_3H_6O_3$) and lactic acid lactate ($C_6H_{10}O_5$). It is obtained by the lactic fermentation of sugars or is prepared synthetically
<i>Note:</i>	
Lactic acid is hygroscopic and when concentrated by boiling, it condenses to form lactic acid lactate, which on dilution and heating hydrolyzes to lactic acid	

Identification

- A. Positive test for lactate

Purity

Sulphated ash	Not more than 0,1 %
Chloride	Not more than 0,2 %
Sulphate	Not more than 0,25 %
Iron	Not more than 10 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg

Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

Note:

This specification refers to a 80 % aqueous solution; for weaker aqueous solutions, calculate values corresponding to their lactic acid content

E 280 PROPIONIC ACID**Definition**

<i>Chemical name</i>	Propionic acid Propanoic acid
<i>Einecs</i>	201-176-3
<i>Chemical formula</i>	$C_3H_6O_2$
<i>Molecular weight</i>	74,08
<i>Assay</i>	Content not less than 99,5 %
<i>Description</i>	Colourless or slightly yellowish, oily liquid with a slightly pungent odour

Identification

A. Melting point	-22 °C
B. Distillation range	138,5 °C to 142,5 °C

Purity

Non-volatile residue	Not more than 0,01 % when dried at 140 °C to constant weight
Aldehydes	Not more than 0,1 % expressed as formaldehyde
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 281 SODIUM PROPIONATE**Definition**

<i>Chemical name</i>	Sodium propionate Sodium propanoate
<i>Einecs</i>	205-290-4
<i>Chemical formula</i>	$C_3H_5O_2Na$
<i>Molecular weight</i>	96,06
<i>Assay</i>	Content not less than 99 % after drying for two hours at 105 °C
<i>Description</i>	White crystalline hygroscopic powder, or a fine white powder

Identification

A. Positive tests for propionate and for sodium

B. pH of a 10 % aqueous solution

Not less than 7,5 and not more than 10,5

Purity

Loss on drying

Not more than 4 % determined by drying for two hours at 105 °C

Water insolubles

Not more than 0,1 %

Iron

Not more than 50 mg/kg

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 282 CALCIUM PROPIONATE**Definition***Chemical name*

Calcium propionate

Einecs

223-795-8

Chemical formula $C_6H_{10}O_4Ca$ *Molecular weight*

186,22

Assay

Content not less than 99 %, after drying for two hours at 105 °C

Description

White crystalline powder

Identification

A. Positive tests for propionate and for calcium

B. pH of a 10 % aqueous solution

Between 6,0 and 9,0

Purity

Loss on drying

Not more than 4 %, determined by drying for two hours at 105 °C

Water insolubles

Not more than 0,3 %

Iron

Not more than 50 mg/kg

Fluoride

Not more than 10 mg/kg

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 283 POTASSIUM PROPIONATE

Definition

<i>Chemical name</i>	Potassium propionate Potassium propanoate
<i>Einecs</i>	206-323-5
<i>Chemical formula</i>	$C_3H_5KO_2$
<i>Molecular weight</i>	112,17
<i>Assay</i>	Content not less than 99 % after drying for two hours at 105 °C
<i>Description</i>	White crystalline powder

Identification

- A. Positive tests for propionate and for potassium

Purity

Loss on drying	Not more than 4 %, determined by drying for two hours at 105 °C
Water-insoluble substances	Not more than 0,3 %
Iron	Not more than 30 mg/kg
Fluoride	Not more than 10 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 284 BORIC ACID

Synonyms

Boracic acid
Orthoboric acid
Borofax

Definition

<i>Einecs</i>	233-139-2
<i>Chemical formula</i>	H_3BO_3
<i>Molecular weight</i>	61,84
<i>Assay</i>	Content not less than 99,5 %
<i>Description</i>	Colourless, odourless, transparent crystals or white granules or powder; slightly unctuous to the touch; occurs in nature as the mineral sassolite

Identification

- | | |
|-----------------------------------|-------------------------|
| A. Melting point | At approximately 171 °C |
| B. Burns with a nice green flame | |
| C. pH of a 3,3 % aqueous solution | Between 3,8 and 4,8 |

Purity

Peroxides	No colour develops with added KI-solution
Arsenic	Not more than 1 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 285 SODIUM TETRABORATE (BORAX)**Synonyms**

Sodium borate

Definition*Chemical name*

Sodium tetraborate
Sodium baborate
Sodium pyroborate
Anhydrous tetraborate

Einecs

215-540-4

Chemical formula

$\text{Na}_2\text{B}_4\text{O}_7$
 $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$

Molecular weight

201,27

Description

Powder or glass-like plates becoming opaque on exposure to air; slowly soluble in water

Identification

A. Melting range

Between 171°C and 175°C with decomposition

Purity

Peroxides	No colour develops with added KI-solution
Arsenic	Not more than 1 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 290 CARBON DIOXIDE**Synonyms**

Carbonic acid gas
Dry ice (solid form)
Carbonic anhydride

Definition*Chemical name*

Carbon dioxide

Einecs

204-696-9

<i>Chemical formula</i>	CO ₂
<i>Molecular weight</i>	44,01
<i>Assay</i>	Content not less than 99 % v/v on the gaseous basis
<i>Description</i>	A colourless gas under normal environmental conditions with a slight pungent odour. Commercial carbon dioxide is shipped and handled as a liquid in pressurized cylinders or bulk storage systems, or in compressed solid blocks of 'dry ice'. Solid (dry ice) forms usually contain added substances, such as propylene glycol or mineral oil, as binders
Identification	
A. Precipitation (Precipitate formation)	When a stream of the sample is passed through a solution of barium hydroxide, a white precipitate is produced which dissolves with effervescence in dilute acetic acid
Purity	
Acidity	915 ml of gas bubbled through 50 ml of freshly boiled water must not render the latter more acid to methylorange than is 50 ml freshly boiled water to which has been added 1 ml of hydrochloric acid (0,01 N)
Reducing substances, hydrogen phosphide and sulphide	915 ml of gas bubbled through 25 ml of ammoniacal silver nitrate reagent to which has been added 3 ml of ammonia must not cause clouding or blackening of this solution
Carbon monoxide	Not more than 10 µl/l
Oil content	Not more than 0,1 mg/l

E 300 ASCORBIC ACID

Definition	
<i>Chemical name</i>	L-ascorbic acid Ascorbic acid 2,3-Didehydro-L-threo-hexono-1,4-lactone 3-Keto-L-gulofuranolactone
Einecs	200-066-2
<i>Chemical formula</i>	C ₆ H ₈ O ₆
<i>Molecular weight</i>	176,13
<i>Assay</i>	Ascorbic acid, after drying in a vacuum desiccator over sulphuric acid for 24 hours, contains not less than 99 % of C ₆ H ₈ O ₆
<i>Description</i>	White to pale yellow, odourless crystalline solid
Identification	
A. Melting range	Between 189 °C and 193 °C with decomposition
B. Positive tests for ascorbic acid	
Purity	
Loss on drying	Not more than 0,4 % after drying in a vacuum desiccator over sulphuric acid for 24 hours
Sulphated ash	Not more than 0,1 %

Specific rotation	$[\alpha]_D^{20}$ between +20,5° and +21,5° (10 % w/v aqueous solution)
pH of a 2 % aqueous solution	Between 2,4 and 2,8
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 301 SODIUM ASCORBATE**Definition**

<i>Chemical name</i>	Sodium ascorbate Sodium L-ascorbate 2,3-Didehydro-L-threo-hexono-1,4-lactone sodium enolate 3-Keto-L-gulofurano-lactone sodium enolate
<i>Einecs</i>	205-126-1
<i>Chemical formula</i>	$C_6H_7O_6Na$
<i>Molecular weight</i>	198,11
<i>Assay</i>	Sodium ascorbate, after drying in a vacuum desiccator over sulphuric acid for 24 hours, contains not less than 99 % of $C_6H_7O_6Na$
<i>Description</i>	White or almost white, odourless crystalline solid which darkens on exposure to light

Identification

- A. Positive tests for ascorbate and for sodium

Purity

Loss on drying	Not more than 0,25 % after drying in a vacuum desiccator over sulphuric acid for 24 hours
Specific rotation	$[\alpha]_D^{20}$ between +103° and +106° (10 % w/v aqueous solution)
pH of 10 % aqueous solution	Between 6,5 and 8,0
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 302 CALCIUM ASCORBATE**Definition**

<i>Chemical name</i>	Calcium ascorbate dihydrate Calcium salt of 2,3-didehydro-L-threo-hexono-1,4-lactone dihydrate
----------------------	---

Einecs	227-261-5
<i>Chemical formula</i>	$C_{12}H_{14}O_{12}Ca \cdot 2H_2O$
<i>Molecular weight</i>	426,35
<i>Assay</i>	Content not less than 98 % on a volatile matter-free basis
<i>Description</i>	White to slightly pale greyish-yellow odourless crystalline powder
Identification	
A. Positive tests for ascorbate and for calcium	
Purity	
Fluoride	Not more than 10 mg/kg (expressed as fluorine)
Specific rotation	$[\alpha]_D^{20}$ between $+95^\circ$ and $+97^\circ$ (5 % w/v aqueous solution)
pH of 10 % aqueous solution	Between 6,0 and 7,5
Volatile matter	Not more than 0,3 % determined by drying at room temperature for 24 hours in a desiccator containing sulphuric acid or phosphorus pentoxide
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 304 (i) ASCORBYL PALMITATE

Definition	
<i>Chemical name</i>	Ascorbyl palmitate L-ascorbyl palmitate 2,3-didehydro-L-threo-hexono-1,4-lactone-6-palmitate 6-palmitoyl-3-keto-L-gulofuranolactone
Einecs	205-305-4
<i>Chemical formula</i>	$C_{22}H_{38}O_7$
<i>Molecular weight</i>	414,55
<i>Assay</i>	Content not less than 98 % on the dried basis
<i>Description</i>	White or yellowish-white solid with a citrus-like odour
Identification	
A. Melting range	Between 107°C and 117°C
Purity	
Loss on drying	Not more than 2,0 % after drying in a vacuum oven at 56°C and 60°C for one hour
Sulphated ash	Not more than 0,1 %

Specific rotation	$[\alpha]_D^{20}$ between +21° and +24° (5 % w/v in methanol solution)
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 304 (ii) ASCORBYL STEARATE**Definition**

<i>Chemical name</i>	Ascorbyl stearate L-ascorbyl stearate 2,3-didehydro-L-threo-hexono-1,4-lactone-6-stearate 6-stearoyl-3-keto-L-gulofuranolactone
<i>Einecs</i>	246-944-9
<i>Chemical formula</i>	$C_{24}H_{42}O_7$
<i>Molecular weight</i>	442,6
<i>Assay</i>	Content not less than 98 %
<i>Description</i>	White or yellowish, white solid with a citrus-like odour

Identification

A. Melting point	About 116°C
------------------	-------------

Purity

Loss on drying	Not more than 2,0 % after drying in a vacuum oven at 56°C to 60°C for one hour
Sulphated ash	Not more than 0,1 %
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 306 TOCOPHEROL-RICH EXTRACT**Definition**

	Product obtained by the vacuum steam distillation of edible vegetable oil products, comprising concentrated tocopherols and tocotrienols Contains tocopherols such as d- α -, d- β -, d- γ - and d- ζ -tocopherols
<i>Molecular weight</i>	430,71 (d- α -tocopherol)
<i>Assay</i>	Content not less than 34 % of total tocopherols
<i>Description</i>	Brownish red to red, clear, viscous oil having a mild, characteristic odour and taste. May show a slight separation of wax-like constituents in microcrystalline form

Identification

A. By suitable gas liquid chromatographic method

B. Solubility tests

Insoluble in water. Soluble in ethanol. Miscible in ether

Purity

Sulphated ash

Not more than 0,1 %

Specific rotation

 $[\alpha]_D^{20}$ not less than +20°

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 307 ALPHA-TOCOPHEROL**Synonyms**dl- α -Tocopherol**Definition***Chemical name*dl-5,7,8-Trimethyltolcol
dl-2,5,7,8-tetramethyl-2-(4',8',12'-trimethyltridecyl)-6-chromanol*Einecs*

200-412-2

Chemical formula $C_{29}H_{50}O_2$ *Molecular weight*

430,71

Assay

Content not less than 96 %

Description

Slightly yellow to amber, nearly odourless, clear, viscous oil which oxidizes and darkens on exposure to air or light

Identification

A. Solubility tests

Insoluble in water, freely soluble in ethanol, miscible in ether

B. Spectrophotometry

In absolute ethanol the maximum absorption is about 292 nm

Purity

Refractive index

 n_D^{20} 1,503 — 1,507Specific absorption $E_{1\text{ cm}}^{1\%}$ in ethanol $E_{1\text{ cm}}^{1\%}$ (292 nm) 72—76
(0,01 g in 200 ml of absolute ethanol)

Sulphated ash

Not more than 0,1 %

Specific rotation

 $[\alpha]_D^{20}$ 0°±0,05° (1 in 10 solution in chloroform)

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 308 GAMMA-TOCOPHEROL

Synonyms	dl- γ -Tocopherol
Definition	
<i>Chemical name</i>	2,7,8-trimethyl-2-(4',8',12'-trimethyltridecyl)-6-chromanol
<i>Einecs</i>	231-523-4
<i>Chemical formula</i>	$C_{28}H_{48}O_2$
<i>Molecular weight</i>	416,69
<i>Assay</i>	Content not less than 97 %
<i>Description</i>	Clear, viscous, pale yellow oil which oxidizes and darkens on exposure to air or light
Identification	
A. Spectrometry	Maximum absorptions in absolute ethanol at about 298 nm and 257 nm
Purity	
Specific absorption $E_{1\text{ cm}}^{1\%}$ in ethanol	$E_{1\text{ cm}}^{1\%}$ (298 nm) between 91 and 97 $E_{1\text{ cm}}^{1\%}$ (257 nm) between 5,0 and 8,0
Refractive index	n_D^{20} 1,503—1,507
Sulphated ash	Not more than 0,1 %
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 309 DELTA-TOCOPHEROL

Definition	
<i>Chemical name</i>	2,8-dimethyl-2-(4',8',12'-trimethyltridecyl)-6-chromanol
<i>Einecs</i>	204-299-0
<i>Chemical formula</i>	$C_{27}H_{46}O_2$
<i>Molecular weight</i>	402,7
<i>Assay</i>	Content not less than 97 %
<i>Description</i>	Clear, viscous, pale yellowish or orange oil which oxidizes and darkens on exposure to air or light
Identification	
A. Spectrometry	Maximum absorptions in absolute ethanol at about 298 nm and 257 nm

Purity

Specific absorption $E_{1\text{ cm}}^{1\%}$ in ethanol	$E_{1\text{ cm}}^{1\%}$ (298 nm) between 89 and 95 $E_{1\text{ cm}}^{1\%}$ (257 nm) between 3,0 and 6,0
Refractive index	n_D^{20} 1,500—1,504
Sulphated ash	Not more than 0,1 %
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 310 PROPYL GALLATE**Definition**

<i>Chemical name</i>	Propyl gallate Propyl ester of gallic acid n-propyl ester of 3,4,5-trihydroxybenzoic acid
<i>Einecs</i>	204-498-2
<i>Chemical formula</i>	$C_{10}H_{12}O_5$
<i>Molecular weight</i>	212,20
<i>Assay</i>	Content not less than 98 % on the anhydrous basis
<i>Description</i>	White to creamy-white, crystalline, odourless solid

Identification

A. Solubility tests	Slightly soluble in water, freely soluble in ethanol, ether and propane-1,2-diol
B. Melting range	Between 146°C and 150°C after drying at 110°C for four hours

Purity

Loss on drying	Not more than 1,0 % (110°C, four hours)
Sulphated ash	Not more than 0,1 %
Free acid	Not more than 0,5 % (as gallic acid)
Chlorinated organic compound	Not more than 100 mg/kg (as Cl)
Specific absorption $E_{1\text{ cm}}^{1\%}$ in ethanol	$E_{1\text{ cm}}^{1\%}$ (275 nm) not less than 485 and not more than 520
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 311 OCTYL GALLATE

Definition

<i>Chemical name</i>	Octyl gallate Octyl ester of gallic acid n-octyl ester of 3,4,5-trihydroxybenzoic acid
<i>Einecs</i>	213-853-0
<i>Chemical formula</i>	$C_{15}H_{22}O_5$
<i>Molecular weight</i>	282,34
<i>Assay</i>	Content not less than 98 % after drying at 90 °C for six hours
<i>Description</i>	White to creamy-white odourless solid

Identification

A. Solubility tests	Insoluble in water, freely soluble in ethanol, ether and propane-1,2-diol
B. Melting range	Between 99 °C and 102 °C after drying at 90 °C for six hours

Purity

Loss on drying	Not more than 0,5 % (90 °C, six hours)
Sulphated ash	Not more than 0,05 %
Free acid	Not more than 0,5 % (as gallic acid)
Chlorinated organic compound	Not more than 100 mg/kg (as C1)
Specific absorption $E_{1\text{cm}}^{1\%}$ in ethanol	$E_{1\text{cm}}^{1\%}$ (275 nm) not less than 375 and not more than 390
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 312 DODECYL GALLATE

Synonyms

Lauryl gallate

Definition

<i>Chemical name</i>	Dodecyl gallate n-dodecyl (or lauryl) ester of 3,4,5-trihydroxybenzoic acid Dodecyl ester of gallic acid
<i>Einecs</i>	214-620-6
<i>Chemical formula</i>	$C_{19}H_{30}O_5$
<i>Molecular weight</i>	338,45
<i>Assay</i>	Content not less than 98 % after drying at 90 °C for six hours
<i>Description</i>	White or creamy-white odourless solid

Identification

- | | |
|---------------------|--|
| A. Solubility tests | Insoluble in water, freely soluble in ethanol and ether |
| B. Melting range | Between 95°C and 98°C after drying at 90°C for six hours |

Purity

- | | |
|--|--|
| Loss on drying | Not more than 0,5 % (90°C, six hours) |
| Sulphated ash | Not more than 0,05 % |
| Free acid | Not more than 0,5 % (as gallic acid) |
| Chlorinated organic compound | Not more than 100 mg/kg (as Cl) |
| Specific absorption $E_{1\text{ cm}}^{1\%}$ in ethanol | $E_{1\text{ cm}}^{1\%}$ (275 nm) not less than 300 and not more than 325 |
| Arsenic | Not more than 3 mg/kg |
| Lead | Not more than 10 mg/kg |
| Mercury | Not more than 1 mg/kg |
| Heavy metals (as Pb) | Not more than 30 mg/kg |

E 315 ERYTHORBIC ACID**Synonyms**

Isoascorbic acid
D-araboascorbic acid

Definition

Chemical name D-Erythro-hex-2-enoic acid γ -lactone
Isoascorbic acid
D-isoascorbic acid

Einecs 201-928-0

Chemical formula $\text{C}_6\text{H}_8\text{O}_6$

Molecular weight 176,13

Assay Content not less than 98 % on the anhydrous basis

Description White to slightly yellow crystalline solid which darkens gradually on exposure to light

Identification

- | | |
|--|---|
| A. Melting range | About 164°C to 172°C with decomposition |
| B. Positive test for ascorbic acid/colour reaction | |

Purity

- | | |
|----------------|---|
| Loss on drying | Not more than 0,4 % after drying under reduced pressure on silica gel for 3 hours |
| Sulphated ash | Not more than 0,3 % |

Specific rotation	$[\alpha]_D^{25}$ 10 % (w/v) aqueous solution between $-16,5^\circ$ to $-18,0^\circ$
Oxalate	To a solution of 1 g in 10 ml of water add 2 drops of glacial acetic acid and 5 ml of 10 % calcium acetate solution. The solution should remain clear
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 316 SODIUM ERYTHORBATE

Synonyms	Sodium isoascorbate
Definition	
<i>Chemical name</i>	Sodium isoascorbate Sodium D-isoascorbic acid Sodium salt of 2,3-didehydro-D-erythro-hexono-1,4-lactone 3-keto-D-gulofurano-lactone sodium enolate monohydrate
Einecs	228-973-9
<i>Chemical formula</i>	$C_6H_7O_6Na \cdot H_2O$
<i>Molecular weight</i>	216,13
Assay	Content not less than 98 % after drying in a vacuum desiccator over sulphuric acid for 24 hours expressed on the monohydrate basis
<i>Description</i>	White crystalline solid
Identification	
A. Solubility tests	Freely soluble in water, very slightly soluble in ethanol
B. Positive test for ascorbic acid/colour reaction	
C. Positive test for sodium	
Purity	
Loss on drying	Not more than 0,25 % after drying in a vacuum desiccator over sulphuric acid for 24 hours
Specific rotation	$[\alpha]_D^{25}$ 10 % (w/v) aqueous solution between $+95^\circ$ and $+98^\circ$
pH of a 10 % aqueous solution	5,5 to 8,0
Oxalate	To a solution of 1 g in 10 ml of water add 2 drops of glacial acetic acid and 5 ml of 10 % calcium acetate solution. The solution should remain clear
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 320 BUTYLATED HYDROXYANISOLE (BHA)

Synonyms	BHA
Definition	
<i>Chemical name</i>	3-Tertiary-butyl-4-hydroxyanisole A mixture of 2-tertiarybutyl-4-hydroxyanisole and 3-tertiarybutyl-4-hydroxyanisole
Einecs	246-563-8
<i>Chemical formula</i>	C ₁₁ H ₁₆ O ₂
<i>Molecular weight</i>	180,25
<i>Assay</i>	Content not less than 98,5 % of C ₁₁ H ₁₆ O ₂ and not less than 85 % of 3-tertiary-butyl-4-hydroxyanisole isomer
<i>Description</i>	White or slightly yellow crystals or waxy solid with a slight aromatic smell
Identification	
A. Solubility tests	Insoluble in water
B. Melting range	Between 48 °C and 55 °C
Purity	
Sulphated ash	Not more than 0,05 % after calcination at 800 ±25 °C
Phenolic impurities	Not more than 0,5 %
Specific absorption E ₁ ^{1%} _{1 cm} in ethanol	E ₁ ^{1%} _{1 cm} (290 nm) not less than 190 and not more than 210 E ₁ ^{1%} _{1 cm} (228 nm) not less than 326 and not more than 345
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 321 BUTYLATED HYDROXYTOLUENE (BHT)

Synonyms	BHT
Definition	
<i>Chemical name</i>	2,6-Ditertiary-butyl- <i>p</i> -cresol 4-Methyl-2,6-ditertiarybutylphenol
Einecs	204-881-4
<i>Chemical formula</i>	C ₁₅ H ₂₄ O
<i>Molecular weight</i>	220,36
<i>Assay</i>	Content not less than 99 %
<i>Description</i>	White, crystalline or flaked solid, odourless or having a characteristic faint aromatic odour

Identification

- | | |
|-----------------------|--|
| A. Solubility tests | Insoluble in water and propane- 1,2-diol .
Freely soluble in ethanol |
| B. Melting point | At 70 °C |
| C. Absorbance maximum | The absorption in the range 230 to 320 nm of a 2 cm layer of a 1 in 100 000 solution in dehydrated ethanol exhibits a maximum only at 278 nm |

Purity

- | | |
|---|---|
| Sulphated ash | Not more than 0,005 % |
| Phenolic impurities | Not more than 0,5 % |
| Specific absorption $E_{1\text{cm}}^{1\%}$ in ethanol | $E_{1\text{cm}}^{1\%}$ (278 nm) not less than 81 and not more than 88 |
| Arsenic | Not more than 3 mg/kg |
| Lead | Not more than 5 mg/kg |
| Mercury | Not more than 1 mg/kg |
| Heavy metals (as Pb) | Not more than 10 mg/kg |

E 322 LECITHINS**Synonyms**

Phosphatides
Phospholipids

Definition

Lecithins are mixtures or fractions of phosphatides obtained by physical procedures from animal or vegetable foodstuffs; they also include hydrolysed products obtained through the use of harmless and appropriate enzymes. The final product must not show any signs of residual enzyme activity

The lecithins may be slightly bleached in aqueous medium by means of hydrogen peroxide. This oxidation must not chemically modify the lecithin phosphatides

Einecs

232-307-2

Assay

- Lecithins: not less than 60,0 % of substances insoluble in acetone
- Hydrolysed lecithins: not less than 56,0 % of substances insoluble in acetone

Description

- Lecithins: brown liquid or viscous semi-liquid or powder
- Hydrolysed lecithins: light brown to brown viscous liquid or paste

Identification

- | | |
|---|--|
| A. Positive tests for choline, for phosphorus and fatty acids | |
| B. Test for hydrolysed lecithin | |

To a 800 ml beaker add 500 ml of water (30 °C—35 °C). Then slowly add 50 ml of the sample with constant stirring. Hydrolysed lecithin will form a homogeneous emulsion. Non-hydrolysed lecithin will form a distinct mass of about 50 g

Purity

- | | |
|--------------------------|---|
| Loss on drying | Not more than 2,0 % determined by drying at 105 °C for one hour |
| Toluene-insoluble matter | Not more than 0,3 % |
| Acid value | — Lecithins: not more than 35 mg of potassium hydroxide per gram
— Hydrolysed lecithins: not more than 45 mg of potassium hydroxide per gram |

Peroxide value	Equal to or less than 10
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 325 SODIUM LACTATE**Definition**

<i>Chemical name</i>	Sodium lactate Sodium 2-hydroxypropanoate
<i>Einecs</i>	200-772-0
<i>Chemical formula</i>	$C_3H_5NaO_3$
<i>Molecular weight</i>	112,06 (anhydrous)
<i>Assay</i>	Content not less than 57 % and not more than 66 %
<i>Description</i>	Colourless, transparent, liquid Odourless, or with a slight, characteristic odour

Identification

- A. Positive test for lactate
- B. Positive test for potassium

Purity

Acidity	Not more than 0,5 % after drying expressed as lactic acid
pH of a 20 % aqueous solution	6,5 to 7,5
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg
Reducing substances	No reduction of Fehling's solution
<i>Note:</i> This specification refers to a 60 % aqueous solution	

E 326 POTASSIUM LACTATE**Definition**

<i>Chemical name</i>	Potassium lactate Potassium 2-hydroxypropanoate
<i>Einecs</i>	213-631-3

<i>Chemical formula</i>	$C_3H_5O_3K$
<i>Molecular weight</i>	128,17 (anhydrous)
<i>Assay</i>	Content not less than 57 % and not more than 66 %
<i>Description</i>	Slightly viscous, almost odourless clear liquid. Odourless, or with a slight, characteristic odour
Identification	
A. Ignition	Ignite potassium lactate solution to an ash. The ash is alkaline, and an effervescence occurs when acid is added
B. Colour reaction	Overlay 2 ml of potassium lactate solution on 5 ml of a 1 in 100 solution of catechol in sulphuric acid. A deep red colour is produced at the zone of contact
C. Positive tests for potassium and for lactate	
Purity	
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg
Acidity	Dissolve 1 g of potassium lactate solution in 20 ml of water, add 3 drops of phenolphthalein TS and titrate with 0,1 N sodium hydroxide. Not more than 0,2 ml should be required
Reducing substances	Potassium lactate solution shall not cause any reduction of Fehling's solution
<i>Note:</i>	
This specification refers to a 60 % aqueous solution	

E 327 CALCIUM LACTATE

Definition	
<i>Chemical name</i>	Calcium dilactate Calcium dilactate hydrate 2-Hydroxypropanoic acid calcium salt
<i>Einecs</i>	212-406-7
<i>Chemical formula</i>	$(C_3H_5O_2)_2 Ca \cdot nH_2O$ (n = 0—5)
<i>Molecular weight</i>	218,22 (anhydrous)
<i>Assay</i>	Content not less than 98 % on the anhydrous basis
<i>Description</i>	Almost odourless, white crystalline powder or granules
Identification	
A. Positive tests for lactate and for calcium	
B. Solubility tests	Soluble in water and practically insoluble in ethanol

Purity

Loss on drying	Determined by drying at 120°C for four hours: — anhydrous: not more than 3,0 % — with 1 molecule of water: not more than 8,0 % — with 3 molecules of water: not more than 20,0 % — with 4,5 molecules of water: not more than 27,0 %
Acidity	Not more than 0,5 % of the dry matter expressed as lactic acid
Fluoride	Not more than 30 mg/kg (expressed as fluorine)
pH of a 5 % solution	Between 6,0 and 8,0
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg
Reducing substances	No reduction of Fehling's solution

E 330 CITRIC ACID**Definition**

<i>Chemical name</i>	Citric acid 2-Hydroxy-1,2,3-propanetricarboxylic acid β-Hydroxytricarballic acid
Einecs	201-069-1
<i>Chemical formula</i>	(a) $C_6H_8O_7$ (anhydrous) (b) $C_6H_8O_7 \cdot H_2O$ (monohydrate)
<i>Molecular weight</i>	(a) 192,13 (anhydrous) (b) 210,15 (monohydrate)
<i>Assay</i>	Citric acid may be anhydrous or it may contain 1 molecule of water. Citric acid contains not less than 99,5 % of $C_6H_8O_7$, calculated on the anhydrous basis
<i>Description</i>	Citric acid is a white or colourless, odourless, crystalline solid, having a strongly acid taste. The monohydrate effloresces in dry air

Identification

A. Solubility tests	Very soluble in water; freely soluble in ethanol; soluble in ether
---------------------	--

Purity

Water content	Anhydrous citric acid contains not more than 0,5 % water; citric acid monohydrate contains not more than 8,8 % water (Karl Fischer method)
Sulphated ash	Not more than 0,05 % after calcination at $800 \pm 25^\circ\text{C}$
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg

Heavy metals (as Pb)	Not more than 5 mg/kg
Oxalates	Not more than 100 mg/kg, expressed as oxalic acid, after drying
Readily carbonizable substances	Heat 1 g of powdered sample with 10 ml of 98 % minimum sulphuric acid in a water bath at 90°C in the dark for one hour. Not more than a pale brown colour should be produced (Matching Fluid K)

E 331 (i) MONOSODIUM CITRATE

Synonyms	Monosodium citrate Monobasic sodium citrate
Definition	
<i>Chemical name</i>	Monosodium citrate Monosodium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid
<i>Chemical formula</i>	(a) $C_6H_7O_7Na$ (anhydrous) (b) $C_6H_7O_7Na \cdot H_2O$ (monohydrate)
<i>Molecular weight</i>	(a) 214,11 (anhydrous) (b) 232,23 (monohydrate)
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	Crystalline white powder or colourless crystals
Identification	
A. Positive tests for citrate and for sodium	
Purity	
Loss on drying	Determined by drying at 180°C for four hours: — anhydrous: not more than 1,0 % — monohydrate: not more than 8,8 %
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
pH of a 1 % aqueous solution	Between 3,5 and 3,8
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg

E 331 (ii) DISODIUM CITRATE

Synonyms	Disodium citrate Dibasic sodium citrate
Definition	
<i>Chemical name</i>	Disodium citrate Disodium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Disodium salt of citric acid with 1,5 molecules of water

Einecs	205-623-3
<i>Chemical formula</i>	$C_6H_6O_7Na_2 \cdot 1,5H_2O$
<i>Molecular weight</i>	263,11
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	Crystalline white powder or colourless crystals
Identification	
A. Positive tests for citrate and for sodium	
Purity	
Loss on drying	Not more than 13,0 % by drying at 180°C for four hours
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
pH of a 1 % aqueous solution	Between 4,9 and 5,2
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg

E 331 (iii) TRISODIUM CITRATE

Synonyms	Trisodium citrate Tribasic sodium citrate
Definition	
<i>Chemical name</i>	Trisodium citrate Trisodium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Trisodium salt of citric acid, in anhydrous, dihydrate or pentahydrate form
Einecs	200-675-3
<i>Chemical formula</i>	Anhydrous: $C_6H_5O_7Na_3$ Hydrated: $C_6H_5O_7Na_3 \cdot nH_2O$ (n = 2 or 5)
<i>Molecular weight</i>	258,07 (anhydrous)
<i>Assay</i>	Not less than 99 % on the anhydrous basis
<i>Description</i>	Crystalline white powder or colourless crystals
Identification	
A. Positive tests for citrate and for sodium	

Purity

Loss on drying	Determined by drying at 180°C for four hours: — anhydrous: not more than 1,0 % — dihydrate: not more than 13,5 % — pentahydrate: not more than 30,3 %
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
pH of a 5 % aqueous solution	Between 7,5 and 9,0
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg

E 332 (i) MONOPOTASSIUM CITRATE**Synonyms**

Monopotassium citrate
Monobasic potassium citrate

Definition

<i>Chemical name</i>	Monopotassium citrate Monopotassium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Anhydrous monopotassium salt of citric acid
<i>Einecs</i>	212-753-4
<i>Chemical formula</i>	$C_6H_7O_7K$
<i>Molecular weight</i>	230,21
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	White, hygroscopic, granular powder or transparent crystals

Identification

- A. Positive tests for citrate and for potassium

Purity

Loss on drying	Not more than 1,0 % determined by drying at 180°C for four hours
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
pH of a 1 % aqueous solution	Between 3,5 and 3,8
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg

E 332 (ii) TRIPOTASSIUM CITRATE

Synonyms	Tripotassium citrate Tribasic potassium citrate
Definition	
<i>Chemical name</i>	Tripotassium citrate Tripotassium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Monohydrated tripotassium salt of citric acid
Einecs	212-755-5
<i>Chemical formula</i>	$C_6H_5O_7K_3 \cdot H_2O$
<i>Molecular weight</i>	324,42
<i>Assay</i>	Content not less than 99 % on the anhydrous basis
<i>Description</i>	White, hygroscopic, granular powder or transparent crystals
Identification	
A. Positive tests for citrate and for potassium	
Purity	
Loss on drying	Not more than 6,0 % determined by drying at 180 °C for four hours
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
pH of a 5 % aqueous solution	Between 7,5 and 9,0
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg

E 333 (i) MONOCALCIUM CITRATE

Synonyms	Monocalcium citrate Monobasic calcium citrate
Definition	
<i>Chemical name</i>	Monocalcium citrate Monocalcium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Monohydrate monocalcium salt of citric acid
<i>Chemical formula</i>	$(C_6H_7O_7)_2Ca \cdot H_2O$
<i>Molecular weight</i>	440,32
<i>Assay</i>	Content not less than 97,5 % on the anhydrous basis
<i>Description</i>	Fine white powder
Identification	
A. Positive tests for citrate and for calcium	

Purity

Loss on drying	Not more than 7,0 % determined by drying at 180 °C for four hours
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
pH of a 1 % aqueous solution	Between 3,2 and 3,5
Fluoride	Not more than 30 mg/kg (expressed as fluorine)
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg
Carbonates	Dissolving 1 g of calcium citrate in 10 ml 2 N hydrochloric acid must not liberate more than a few isolated bubbles

E 333 (ii) DICALCIUM CITRATE**Synonyms**

Dicalcium citrate
Dibasic calcium citrate

Definition

<i>Chemical name</i>	Dicalcium citrate Dicalcium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Trihydrated dicalcium salt of citric acid
<i>Chemical formula</i>	$(C_6H_7O_7)_2Ca_2 \cdot 3H_2O$
<i>Molecular weight</i>	530,42
<i>Assay</i>	Not less than 97,5 % on the anhydrous basis
<i>Description</i>	Fine white powder

Identification

- A. Positive tests for citrate and for calcium

Purity

Loss on drying	Not more than 20,0 % determined by drying at 180 °C for four hours
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
Fluoride	Not more than 30 mg/kg (expressed as fluorine)
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg
Carbonates	Dissolving 1 g of calcium citrate in 10 ml 2 N hydrochloric acid must not liberate more than a few isolated bubbles

E 333 (iii) TRICALCIUM CITRATE

Synonyms	Tricalcium citrate Tribasic calcium citrate
Definition	
<i>Chemical name</i>	Tricalcium citrate Tricalcium salt of 2-hydroxy-1,2,3-propanetricarboxylic acid Tetrahydrated tricalcium salt of citric acid
Einecs	212-391-7
<i>Chemical formula</i>	$(C_6H_6O_7)_2Ca_3 \cdot 4H_2O$
<i>Molecular weight</i>	570,51
<i>Assay</i>	Not less than 97,5 % on the anhydrous basis
<i>Description</i>	Fine white powder
Identification	
A. Positive tests for citrate and for calcium	
Purity	
Loss on drying	Not more than 14,0 % determined by drying at 180°C for four hours
Oxalates	Not more than 100 mg/kg expressed as oxalic acid, after drying
Fluoride	Not more than 30 mg/kg (expressed as fluorine)
Arsenic	Not more than 1 mg/kg
Lead	Not more than 1 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 5 mg/kg
Carbonates	Dissolving 1 g of calcium citrate in 10 ml 2 N hydrochloric acid must not liberate more than a few isolated bubbles

E 334 L(+)-TARTARIC ACID

Definition	
<i>Chemical name</i>	L-tartaric acid L-2,3-dihydroxybutanedioic acid d- α , β -dihydroxysuccinic acid
Einecs	201-766-0
<i>Chemical formula</i>	$C_4H_6O_6$
<i>Molecular weight</i>	150,09
<i>Assay</i>	Content not less than 99,5 % on the anhydrous basis
<i>Description</i>	Colourless or translucent crystalline solid or white crystalline powder

Identification

- A. Melting range Between 168 °C and 170 °C
- B. Positive test for tartrate

Purity

- Loss on drying Not more than 0,5 % (over P₂O₅, three hours)
- Sulphated ash Not more than 1 000 mg/kg after calcination at 800 ± 25 °C
- Specific optical rotation of a 20 % w/v aqueous solution $[\alpha]_D^{20}$ between +11,5° and +13,5°
- Lead Not more than 5 mg/kg
- Mercury Not more than 1 mg/kg
- Heavy metals (as Pb) Not more than 10 mg/kg
- Oxalates Not more than 100 mg/kg expressed as oxalic acid, after drying

E 335 (i) MONOSODIUM TARTRATE**Synonyms**

Monosodium salt of L-(+)-tartaric acid

Definition

- Chemical name* Monosodium salt of L-2,3-dihydroxybutanedioic acid
Monohydrated monosodium salt of L-(+)-tartaric acid
- Chemical formula* C₄H₅O₆Na·H₂O
- Molecular weight* 194,05
- Assay* Content not less than 99 % on the anhydrous basis
- Description* Transparent colourless crystals

Identification

- A. Positive tests for tartrate and for sodium

Purity

- Loss on drying Not more than 10,0 % determined by drying at 105 °C for four hours
- Oxalates Not more than 100 mg/kg expressed as oxalic acid, after drying
- Arsenic Not more than 3 mg/kg
- Lead Not more than 5 mg/kg
- Mercury Not more than 1 mg/kg
- Heavy metals (as Pb) Not more than 10 mg/kg

E 335 (ii) DISODIUM TARTRATE

Definition

Chemical name

Disodium L-tartrate
 Disodium (+)-tartrate
 Disodium (+)-2,3-dihydroxybutanedioic acid
 Dihydrated disodium salt of L-(+)-tartaric acid

Einecs

212-773-3

Chemical formula $C_4H_4O_6Na_2 \cdot 2H_2O$ *Molecular weight*

230,8

Assay

Content not less than 99 % on the anhydrous basis

Description

Transparent, colourless crystals

Identification

A. Positive tests for tartrate and for sodium

B. Solubility tests

1 gram is insoluble in 3 ml of water. Insoluble in ethanol

Purity

Loss on drying

Not more than 17,0 % determined by drying at 150 °C for four hours

Oxalates

Not more than 100 mg/kg expressed as oxalic acid, after drying

pH of a 1 % aqueous solution

Between 7,0 and 7,5

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 336 (i) MONOPOTASSIUM TARTRATE

Synonyms

Monobasic potassium tartrate

Definition

Chemical name

Anhydrous monopotassium salt of L-(+)-tartaric acid
 Monopotassium salt of L-2,3-dihydroxybutanedioic acid

Chemical formula $C_4H_5O_6K$ *Molecular weight*

188,16

Assay

Content not less than 98 % on the anhydrous basis

Description

White crystalline or granulated powder

Identification

A. Positive tests for tartrate and for potassium

B. Melting point

230°C

Purity

pH of a 1 % aqueous solution

3,4

Loss on drying

Not more than 1,0 % determined by drying at 105°C for four hours

Oxalates

Not more than 100 mg/kg expressed as oxalic acid, after drying

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 336 (ii) DIPOTASSIUM TARTRATE**Synonyms**

Dibasic potassium tartrate

Definition*Chemical name*Dipotassium salt of L-2,3-dihydroxybutanedioic acid
Dipotassium salt with half a molecule of water of L-(+)-tartaric acid*Einecs*

213-067-8

Chemical formula $C_4H_4O_6K_2 \cdot \frac{1}{2}H_2O$ *Molecular weight*

235,2

Assay

Content not less than 99 % on the anhydrous basis

Description

White crystalline or granulated powder

Identification

A. Positive tests for tartrate and for potassium

Purity

pH of a 1 % aqueous solution

Between 7,0 and 9,0

Loss on drying

Not more than 4,0 % determined by drying at 150°C for four hours

Oxalates

Not more than 100 mg/kg expressed as oxalic acid, after drying

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 337 POTASSIUM SODIUM TARTRATE

Synonyms

Potassium sodium L-(+)-tartrate
Rochelle salt
Seignette salt

Definition

Chemical name

Potassium sodium salt of L-2,3-dihydroxybutanedioic acid
Potassium sodium L-(+)-tartrate

Einecs

206-156-8

Chemical formula $C_4H_4O_6KNa \cdot 4H_2O$ *Molecular weight*

282,23

Assay

Content not less than 99 % on the anhydrous basis

Description

Colourless crystals or white crystalline powder

Identification

A. Positive tests for tartrate, for potassium and for sodium

B. Solubility tests

1 gram is soluble in 1 ml of water, insoluble in ethanol

C. Melting range

Between 70 and 80 °C

Purity

Loss on drying

Not more than 26,0 % and not less than 21,0 % determined by drying at 150 °C for three hours

Oxalates

Not more than 100 mg/kg expressed as oxalic acid, after drying

pH of 1 % aqueous solution

Between 6,5 and 8,5

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 338 PHOSPHORIC ACID

Synonyms

Orthophosphoric acid
Monophosphoric acid

Definition

Chemical name

Phosphoric acid

Einecs

231-633-2

Chemical formula H_3PO_4 *Molecular weight*

98,00

Assay

Content not less than 71 % and not more than 83 %

Description

Clear, colourless, viscous liquid

Identification

- A. Positive tests for acid and for phosphate

Purity

Volatile acids	Not more than 10 mg/kg (as acetic acid)
Chlorides	Not more than 200 mg/kg (expressed as chlorine)
Nitrates	Not more than 5 mg/kg (as NaNO_3)
Sulphates	Not more than 1 500 mg/kg (as CaSO_4)
Fluoride	Not more than 10 mg/kg (expressed as fluorine)
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

Note:

This specification refers to a 75 % aqueous solution

E 339 (i) MONOSODIUM PHOSPHATE**Synonyms**

Monosodium monophosphate
Acid monosodium monophosphate
Monosodium orthophosphate
Monobasic sodium phosphate

Definition*Chemical name*

Sodium dihydrogen monophosphate

Einecs

231-449-2

Chemical formula

Anhydrous: NaH_2PO_4
Monohydrate: $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$
Dihydrate: $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$

Molecular weight

Anhydrous: 119,98
Monohydrate: 138,00
Dihydrate: 156,01

Assay

After drying at 6,0°C for one hour and then at 105°C for four hours, contains not less than 97 % of NaH_2PO_4

Description

A white odourless, slightly deliquescent powder, crystals or granules

Identification

- A. Positive tests for sodium and for phosphate
- B. Solubility tests
- C. P_2O_5 content

Freely soluble in water. Insoluble in ethanol, ether or chloroform

Between 58,0 % and 60,0 %

Purity

Loss on drying	The anhydrous salt loses no more than 2,0 %, the monohydrate no more than 15,0 %, and the dihydrate no more than 25 % when dried first at 60°C for one hour, then at 105°C for four hours
Water-insoluble substances	Not more than 0,2 % on the anhydrous basis
Fluoride	Not more than 10 mg/kg (expressed as fluorine)
pH of a 1 % aqueous solution	Between 4,1 and 5,0
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 339 (ii) DISODIUM PHOSPHATE**Synonyms**

Disodium monophosphate
 Secondary sodium phosphate
 Disodium orthophosphate
 Acid disodium phosphate

Definition

<i>Chemical name</i>	Disodium hydrogen monophosphate Disodium hydrogen orthophosphate
Einecs	231-448-7
<i>Chemical formula</i>	Anhydrous: Na_2HPO_4 Hydrated: $\text{Na}_2\text{HPO}_4 \cdot n\text{H}_2\text{O}$ (n = 2, 7 or 12)
<i>Molecular weight</i>	141,98 (anhydrous)
<i>Assay</i>	After drying at 40°C for three hours and subsequently at 105°C for five hours, contains not less than 98 % of Na_2HPO_4
<i>Description</i>	Anhydrous disodium hydrogen phosphate is a white, hygroscopic, odourless powder. Hydrated forms available include the dihydrate: a white crystalline, odourless solid; the heptahydrate: white, odourless, efflorescent crystals or granular powder; and the dodecahydrate: white, efflorescent, odourless powder or crystals

Identification

A. Positive tests for sodium and for phosphate	
B. Solubility tests	Freely soluble in water. Insoluble in ethanol
C. P_2O_5 content	Between 49 % and 51 % (anhydrous)

Purity

Loss on drying	When dried at 40°C for three hours and then at 105°C for five hours, the losses in weight are as follows: anhydrous not more than 5,0 %, dihydrate not more than 22,0 %, heptahydrate not more than 50,0 %, dodecahydrate not more than 61,0 %
----------------	--

Water-insoluble substances	Not more than 0,2 % on the anhydrous basis
Fluoride	Not more than 10 mg/kg (expressed as fluorine)
pH of a 1,0 % aqueous solution	Between 8,4 and 9,6
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 339 (iii) TRISODIUM PHOSPHATE

Synonyms

Sodium phosphate
Tribasic sodium phosphate
Trisodium orthophosphate

Definition

Chemical name

Trisodium monophosphate
Trisodium phosphate
Trisodium orthophosphate

Einecs

231-509-8

Chemical formula

Anhydrous: Na_3PO_4
Hydrated: $\text{Na}_3\text{PO}_4 \cdot n\text{H}_2\text{O}$ ($n = 0,5, 1 \text{ or } 12$)

Molecular weight

163,94 (anhydrous)

Assay

Sodium phosphate anhydrous, and also the hemi- and monohydrates, contains not less than 97,0 % of Na_3PO_4 , calculated on the dried basis. Sodium phosphate dodecahydrate contains not less than 92,0 % of Na_3PO_4 , calculated on the ignited basis

Description

White odourless crystals, granules or a crystalline powder. Hydrated forms available include hemi- and monohydrates, hexahydrate, octahydrate, decahydrate and dodecahydrate. The dodecahydrate contains $\frac{1}{4}$ molecule of sodium hydroxide

Identification

A. Positive tests for sodium and for phosphate

B. Solubility tests

Freely soluble in water. Insoluble in ethanol

C. P_2O_5 content

Between 40,5 % and 43,5 % (anhydrous)

Purity

Loss on ignition

When dried at 120°C for two hours and then ignited at about 800°C for 30 minutes, the losses in weight are as follows: anhydrous not more than 2,0 %, monohydrate: not more than 11,0 %, dodecahydrate: between 45,0 % and 58,0 %

Water-insoluble substances

Not more than 0,2 % on the anhydrous basis

Fluoride

Not more than 10 mg/kg (expressed as fluorine)

pH of a 1,0 % aqueous solution	Between 11,5 and 12,5
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 340 (i) MONOPOTASSIUM PHOSPHATE

Synonyms

Monobasic potassium phosphate
 Monopotassium monophosphate
 Potassium acid phosphate
 Potassium orthophosphate

Definition

Chemical name

Potassium dihydrogen phosphate
 Monopotassium dihydrogen orthophosphate
 Monopotassium dihydrogen monophosphate

Einecs

231-913-4

Chemical formula KH_2PO_4 *Molecular weight*

136,09

Assay

Content not less than 98,0 % after drying at 105 °C for four hours

Description

Odourless, colourless crystals or white granular or crystalline powder, hygroscopic

Identification

A. Positive tests for potassium and for phosphate

B. Solubility tests

Freely soluble in water. Insoluble in ethanol

C. P_2O_5 content

Between 51,0 % and 53,0 %

Purity

Loss on drying

Not more than 2,0 % determined by drying at 105 °C for four hours

Water-insoluble substances

Not more than 0,2 % on the anhydrous basis

Fluoride

Not more than 10 mg/kg (expressed as fluorine)

pH of a 1 % aqueous solution

Between 4,2 and 4,8

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 340 (ii) DIPOTASSIUM PHOSPHATE

Synonyms

Dipotassium monophosphate
Secondary potassium phosphate
Dipotassium acid phosphate
Dipotassium orthophosphate
Dibasic potassium phosphate

Definition

Chemical name

Dipotassium hydrogen monophosphate
Dipotassium hydrogen phosphate
Dipotassium hydrogen orthophosphate

Einecs

231-834-5

Chemical formula K_2HPO_4 *Molecular weight*

174,18

Assay

Content not less than 98 % after drying at 105 °C for four hours

Description

Colourless or white granular powder, crystals or masses; deliquescent substance

Identification

A. Positive tests for potassium and for phosphate

B. Solubility tests

Freely soluble in water. Insoluble in ethanol

C. P_2O_5 content

Between 40,3 % and 41,5 %

Purity

Loss on drying

Not more than 2,0 % determined by drying at 105 °C for four hours

Water-insoluble substances

Not more than 0,2 % on the anhydrous basis

Fluoride

Not more than 10 mg/kg (expressed as fluorine)

pH of a 1 % aqueous solution

Between 8,7 and 9,4

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 340 (iii) TRIPOTASSIUM PHOSPHATE

Synonyms

Potassium phosphate
Tribasic potassium phosphate
Tripotassium orthophosphate

Definition

Chemical name

Tripotassium monophosphate
Tripotassium phosphate
Tripotassium orthophosphate

Einecs	231-907-1
<i>Chemical formula</i>	Anhydrous: K_3PO_4 Hydrated: $K_3PO_4 \cdot nH_2O$ ($n = 1$ or 3)
<i>Molecular weight</i>	212,27 (anhydrous)
<i>Assay</i>	Content not less than 97 % calculated on the ignited basis
<i>Description</i>	Colourless or white, odourless hygroscopic crystals or granules. Hydrated forms available include the monohydrate and trihydrate
Identification	
A. Positive tests for potassium and for phosphate	
B. Solubility tests	Freely soluble in water. Insoluble in ethanol
C. P_2O_5 content	Between 30,5 % and 33,0 % (anhydrous on ignited basis)
Purity	
Loss on ignition	Anhydrous: not more than 3,0 %; hydrated: not more than 23,0 %. Determined by drying at 105°C for one hour and then ignite at about 800°C \pm 25°C for 30 minutes
Water-insoluble substances	Not more than 0,2 % on the anhydrous basis
Fluoride	Not more than 10 mg/kg (expressed as fluorine)
pH of a 1 % aqueous solution	Between 11,5 and 12,3
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 341 (i) MONOCALCIUM PHOSPHATE

Synonyms	Monobasic calcium phosphate Monocalcium orthophosphate
Definition	
<i>Chemical name</i>	Calcium dihydrogen phosphate
Einecs	231-837-1
<i>Chemical formula</i>	Anhydrous: $Ca(H_2PO_4)_2$ Monohydrate: $Ca(H_2PO_4)_2 \cdot H_2O$
<i>Molecular weight</i>	234,05 (anhydrous) 252,08 (monohydrate)
<i>Assay</i>	Content not less than 95 % on the dried basis
<i>Description</i>	Granular powder or white, deliquescent crystals or granules

Identification

A. Positive tests for calcium and for phosphate

B. P_2O_5 content

Between 55,5 % and 61,1 % (anhydrous)

C. CaO content

Between 23,0 % and 27,5 % (anhydrous)

Between 19,0 % and 24,8 % (monohydrate)

Purity

Loss on drying

Not less than 14 % determined by drying at 105°C for four hours (anhydrous)

Not more than 17,5 % determined by drying at 60°C for one hour, then at 105°C for four hours (monohydrate)

Loss on ignition

Not more than 17,5 % after ignition at 800°C±25°C for 30 minutes (anhydrous)

Not more than 25,0 % determined by drying at 105°C for one hour, then ignite at 800°C ±25°C for 30 minutes (monohydrate)

Fluoride

Not more than 30 mg/kg (expressed as fluorine)

Arsenic

Not more than 3 mg/kg

Lead

Not more than 5 mg/kg

Mercury

Not more than 1 mg/kg

Heavy metals (as Pb)

Not more than 10 mg/kg

E 341 (ii) DICALCIUM PHOSPHATE**Synonyms**Dibasic calcium phosphate
Dicalcium orthophosphate**Definition***Chemical name*Calcium monohydrogen phosphate
Calcium hydrogen orthophosphate
Secondary calcium phosphate*Einecs*

231-826-1

*Chemical formula*Anhydrous: $CaHPO_4$ Dihydrate: $CaHPO_4 \cdot 2H_2O$ *Molecular weight*

136,06 (anhydrous)

172,09 (dihydrate)

*Assay*Dicalcium phosphate, after drying at 200°C for three hours, contains not less than 98 % and not more than the equivalent of 102% of $CaHPO_4$ *Description*

White crystals or granules, granular powder or powder

Identification

A. Positive tests for calcium and for phosphate

B. Solubility tests

Sparingly soluble in water. Insoluble in ethanol

C. P_2O_5 content

Between 50,0 % and 52,5 % (anhydrous)

Purity

Loss on ignition	Not more than 8,5 % (anhydrous), or 26,5 % (dihydrate) after ignition at 800 °C \pm 25 °C for 30 minutes
Fluoride	Not more than 50 mg/kg
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 341 (iii) TRICALCIUM PHOSPHATE**Synonyms**

Calcium phosphate, tribasic
Calcium orthophosphate

Definition*Chemical name*

Tricalcium monophosphate

Einecs

231-840-8

Chemical formula

$\text{Ca}_3(\text{PO}_4)_2$

Molecular weight

310,17

Assay

Not less than 90 % calculated on the ignited basis

Description

A white, odourless and tasteless powder which is stable in air

Identification

A. Positive tests for calcium and for phosphate

B. Solubility tests

Practically insoluble in water; insoluble in ethanol, soluble in dilute hydrochloric and nitric acid

C. P_2O_5 content

Between 38,5 % and 48,0 % (anhydrous)

Purity

Loss on ignition	Not more than 8 % after ignition at 800 °C \pm 25 °C, to constant weight
Fluoride	Not more than 50 mg/kg (expressed as fluorine)
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 385 CALCIUM DISODIUM ETHYLENEDIAMINETETRAACETATE

Synonyms	Calcium disodium EDTA Calcium disodium edetate
Definition	
<i>Chemical name</i>	N,N'-1,2-Ethanediyibis [N-(carboxymethyl)-glycinate] [(4-)-O,O',O ^N ,O ^N]calciate(2)-disodium Calcium disodium ethylenediaminetetra acetate Calcium disodium (ethylenedinitrilo)tetra acetate
Einecs	200-529-9
<i>Chemical formula</i>	C ₁₀ H ₁₂ O ₈ CaN ₂ Na ₂ ·2H ₂ O
<i>Molecular weight</i>	410,31
<i>Assay</i>	Content not less than 97 % on the anhydrous basis
<i>Description</i>	White, odourless crystalline granules or white to nearly white powder, slightly hygroscopic
Identification	
A. Positive tests for sodium and for calcium	
B. Chelating activity to metal ions positive	
C. pH of a 1 % solution between 6,5 and 7,5	
Purity	
Water content	5 to 13 % (Karl Fischer method)
Arsenic	Not more than 3 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg

E 1105 LYSOZYME

Synonyms	Lysozyme hydrochloride Muramidase
Definition	Lysozyme is a linear polypeptide obtained from hens' egg whites consisting of 129 amino acids. It possesses enzymatic activity in its ability to hydrolyse the β(1-4) linkages between N-acetylmuramic acid and N-acetylglucosamine in the outer membranes of bacterial species, in particular gram-positive organisms. Is usually obtained as the hydrochloride
<i>Chemical name</i>	Enzyme Commission (EC) No: 3.2.1.17
Einecs	232-620-4

<i>Molecular weight</i>	About 14 000
<i>Assay</i>	Content not less than 950 mg/g on the anhydrous basis
<i>Description</i>	White, odourless powder having a slightly sweet taste
Identification	
A. Isoelectric point 10,7	
B. pH of a 2 % aqueous solution between 3,0 and 3,6	
C. Absorption maximum of an aqueous solution (25 mg/100 ml) at 281 nm, a minimum at 252 nm	
Purity	
Water content	Not more than 6,0 % (Karl Fischer method) (powder form only)
Residue on ignition	Not more than 1,5 %
Nitrogen	Not less than 16,8 % and not more than 17,8 %
Arsenic	Not more than 1 mg/kg
Lead	Not more than 5 mg/kg
Mercury	Not more than 1 mg/kg
Heavy metals (as Pb)	Not more than 10 mg/kg
Microbiological criteria	
Total bacterial count	Not more than 5×10^4 col/g
<i>Salmonellae</i>	Absent in 25 g
<i>Staphylococcus aureus</i>	Absent in 1 g
<i>Escherichia coli</i>	Absent in 1 g